

## **NOVEL-CYTIDINE DEAMINASE**

[001] This application is a continuation-in-part of PCT/JP00/01918, filed March 28, 2000,  
5 and claims priority from Japanese Patent Application No. 11-87192, filed March 29, 1999;  
Japanese Patent Application No. 11/178999, filed June 24, 1999; and Japanese Patent  
Application No. 11/371382, filed December 27, 1999.

### **FIELD OF INVENTION**

[002] The present invention relates to novel proteins having a cytidine deaminase activity;  
10 DNAs and fragments thereof (cDNAs, genomic DNAs, and primer DNAs) encoding the  
proteins; expression vectors comprising the DNAs; transformants transformed with the  
expression vectors; antibodies reactive to the proteins or fragments thereof; cells producing  
the antibodies; and methods for identifying substances that regulate production of the  
proteins, transcription of genes encoding the proteins into mRNAs, or enzyme activities of  
15 the proteins.-

### **BACKGROUND**

[003] The germinal center of mammals comprises a highly specialized microenvironment  
required for the final process of maturation towards antigen specific memory cells and long-  
lived plasma cells (Embo J., 16:2996-3006, 199; Semin. Immunol., 4:11-17, 1992). In the  
20 this microenvironment, it is known that two fundamental editings of the immunoglobulin  
genes take place (J. Exp. Med., 173:1165-1175, 1991; Embo. J., 12:4955-4967, 1993; Adv.  
Exp. Med. Biol., 186:145-151, 1985; Nature, 342:929-931, 1989; Cell, 67:1121-1129).

[004] The first fundamental editing is ~~One is the~~ somatic hypermutation (Curr. Opin.  
Immunol., 7:248-254, 1995; Annu. Rev. Immunol., 14:441-457, 1996; Science, 244:1152-  
25 1157, 1989), a phenomenon in which extensive point mutation in the exons of ~~exon~~ genes  
encoding variable regions of immunoglobulins occurs. Accumulation of ~~the~~ point mutations  
leads to selection of B cells expressing high affinity immunoglobulins on their cell surface,  
accompanied by the affinity maturation of antibodies (Embo. J., 4:345-350, 1985; Proc. Natl.

Acad. Sci. USA, 85:8206-8210, 1988). As ~~the~~ a result, immunoglobulin genes are edited as new functional genes.

[005] Another fundamental editing process is the class switch recombination (CSR). In ~~the recombination~~ CSR, effector functions of antibodies, such as complement fixation, are  
5 selected by exchanging exons encoding constant regions of immunoglobulin heavy chains (Curr. Top. Microbiol. Immunol., 217:151-169, 1996; Annu. Rev. Immunol., 8:717-735, 1990).

[006] These two types of genetic editing are very important for effective humoral immunoreaction to eliminate harmful microbes. The molecular mechanisms of the genetic  
10 phenomena have not yet been elucidated despite ~~the extensive studies~~ study for several decades.

[007] The present inventors isolated a mouse B cell clone, CH12F3-2, as a research tool to elucidate the molecular mechanism of class switch recombination of immunoglobulin. In ~~the~~ this B cell line, class switch recombination (CSR) from IgM to IgA begins several hours after  
15 stimulation with IL-4, TGF- $\beta$ , and CD40L; ~~and finally~~ ultimately, over 80% of the cells become IgA positive (Immunity, 9:1-10, 1998; Curr. Biol., 8:227-230, 1998; Int. Immunol., 8:193-201, 1996).

[008] Using the mouse B cell clone CH12F3-2, the present inventors ~~had previously~~ reported that the breakpoints of CSR distribute not only in the switch region (or "S region"),  
20 ~~characterized~~ dist ~~ie~~ by repeated sequences, but also in ~~the~~ neighboring sequences (Curr. Biol., 8:227-230, 1998). However, the breakpoints were rarely seen in I exon and C exon, which are located ~~ing at~~ upstream and downstream of the S region, respectively. Also, according to ~~the~~ accumulated scientific evidence, it has been shown that ~~the~~ transcription of I exon and C exon and ~~the~~ splicing of the transcripts are essential for CSR (Cell, 73:1155-1164, 1993;  
25 Science, 259:984-987, 1993; Proc. Natl. Acad. Sci, USA, 90:3705-3709, 1993; Cell, 81:833-836, 1995).

[009] This ~~namely~~ suggests that the transcripts are involved in CSR either directly or indirectly. Accordingly, the present inventors propose a theory that class switch is initiated by the recognition of DNA-RNA complex structure and not by the recognition of nucleotide  
30 sequences of the switch region. This idea is further fortified by the fact that even when the ~~in the case that~~ Sa region is substituted with an S $\alpha$  region or an S $\gamma$  region by introducing a mini-

chromosome ~~into the~~ above-mentioned mouse B cell clone CH12F3-2, CSR in the mini-chromosome efficiently occurs after stimulation with ~~by stimuli of~~ cytokines (Immunity, 9:1-10, 1998).

[0010] In plants and ~~P~~protozoa, RNA editing, another type of genetic editing, is widely used  
5 as a mean for producing functional genes from a limited genome (Cell, 81:833-836, 1995; Cell, 81:837-840, 1995). mRNA editing of many molecules such as the mRNA ~~of~~ for apolipoprotein B (apoB), AMPA receptors, Wilmstumor-1,  $\alpha$ -galactosidase and neurofibromatosis type-1, and tRNA-Asp, have been reported (Trends Genet., 12:418-424, 1996; Curr. Opin. Genet. Dev., 6:221-231, 1996). Although the molecular mechanism of  
10 mammalian RNA editing has not yet been elucidated, one performed by APOBEC-1 (apolipoprotein B mRNA editing enzyme, catalytic polypeptide-1) is becoming ~~becomes~~ understood by degrees (Science, 260:1816-1819, 1993; J. Biol. Chem., 268:20709-20712, 1993).

[0011] In apoB RNA editing, the first base C (cytosine) of codon CAA, which encodes  
15 glutamine, is converted to U (uridine), which alters the codon to UAA. As a ~~the~~ result, an in-frame stop codon is made in the apoB mRNA (J. Cell., 81:187-195, 1995; J. Cell., 50:831-840, 1987; Science, 238:363-266, 1987). apoB-48 and apoB-100 are transcripts of edited mRNA and unedited mRNA of apoB, respectively, and these proteins possess totally different physiological functions from ~~or~~ each other (J. Biol. Chem., 271:2353-2356, 1996).

[0012] In ~~the~~ site-specific RNA-editing, auxiliary factors are required (Science, 260:1816-1819, 1993; J. Biol. Chem., 268:20709-20712, 1993). In the absence of auxiliary factors, APOBEC-1 shows only a cytidine deaminase activity, possessing non-specific low affinity to RNA (J. Biol. Chem., 268:20709-20712, 1993; J. Cell., 81:187-195, 1995; J. Biol. Chem., 270:14768-14775, 1995; J. Biol. Chem., 270:14762-14767, 1995). The expression and  
25 activity of the auxiliary factors are found not only in organs with apoB mRNA editing, but also in organs with undetectable levels of APOBEC-1 expression, or organs without apoB mRNA editing (Science, 260:1816-1819, 1993; J. Biol. Chem., 268:20709-20712, 1993; Nucleic Acids Res., 22:1874-1879, 1994; Proc Natl. Acad. Sci, USA, 91:8522-8526, 1994; J. Biol. Chem., 269:21725-21734, 1994).

[0013] The unexpected expression of the auxiliary factors involved in apoB mRNA editing  
30 suggests that the auxiliary factors may be involved in more general cellular functions or other

yet unknown RNA editing. Since ~~there are possibilities~~ the possibility exists that CSR and hypermutation, which are involved in genetic editing of relating to immunoglobulin genes, may be accomplished by RNA editing, it ~~is~~ would be very interesting to elucidate whether RNA editing takes place or not in the genetic editing of immunoglobulin genes as mentioned above.

## SUMMARY

[0014] The present invention provides AID (Activation-Induced cytidine Deaminase), a novel cytidine deaminase ~~having that is~~ structurally related relationship to APOBEC-1, ~~one of an~~ RNA editing enzyme, and is involved in RNA editing in germinal center B cells, where genetic editing of immunoglobulin genes occurs, and a DNA encoding the new enzyme.

[0015] The present inventors intensively searched for novel genes involved in class switch recombination (CSR), one of the major types of genetic editing of immunoglobulin genes. As a result, by preparing cDNA libraries for the mouse B cell clone CH12F3-2, (in which class switch recombination from IgM to IgA is shown to occur at an extremely high rate upon together with activation of the cells by stimulation with cytokines), with and without stimulation ~~ing~~ with cytokines, and performing subtraction cloning using the libraries, the present inventors found genes encoding mouse- and human-derived novel proteins named AID (Activation-Induced cytidine Deaminase), having a structural relationship to APOBEC-1, one of the RNA editing enzymes, and having a cytidine deaminase activity similar to APOBEC-1.

[0016] The AID protein in the present invention possesses features described below, and is considered to be a very important RNA-modifying deaminase involved in regulating B cell activation, CSR of immunoglobulin genes, somatic hypermutation, and affinity maturation, which ~~all are~~ all involved in genetic editing specific to germinal center function:

(1) The ORF of the cDNA encoding the AID protein comprises 198 amino acids, with a 24kDa calculated molecular weight (mouse: SEQ ID NO:2, and human: SEQ ID NO:8). ~~Mouse~~ The mouse AID protein shows an approximately 28kDa molecular weight by SDS-PAGE.

(2) The amino acid sequence of the AID protein is 34% and 26% identical to APOBEC-1 (apolipoprotein B mRNA editing enzyme, catalytic polypeptide-1) at the amino acid sequence level, for mouse and human derived proteins, respectively.

(3) The AID protein has a cytidine/deoxycytidine deaminase motif, which is the active center of the deaminase activity ~~which~~ and is conserved in amino acid sequences of proteins belonging to the cytosine nucleoside/nucleotide deaminase family.

(4) The ~~c~~Cytidine deaminase motif of the AID protein is allied with an RNA editing deaminase subgroup.

(5) The AID protein has a ~~L~~leucine-rich region considered to be important in protein-protein interaction, similar to APOBEC-1. Four leucines in this leucine-rich region of the AID protein are conserved in the leucine-rich region of APOBEC-1 in rabbit, rat, mouse and human.

(6) In the primary structure of the AID protein, all of the amino acid residues reported to be necessary for APOBEC-1 to bind RNA (Phe66, Phe87, His61, Glu63 and Cys93) are conserved.

(7) The AID protein has a pseudoactive site domain in its C terminal for forming homodimers, similar to APOBEC-1 and ECDDA, an *E. coli* derived cytidine deaminase. There ~~is a~~ ~~are possibilities~~ possibility that the AID protein forms homodimers, or associates with other auxiliary proteins.

(8) The AID protein shows a concentration-dependent cytidine deaminase activity. The activity can be inhibited dose dependently by tetrahydrouridine (THU), a specific inhibitor of cytidine deaminase. Also, a zinc chelator, 1,10-o-phenanthroline, inhibits the cytidine deaminase activity of the AID protein while 1,7-o-phenanthroline, the inactive isomer, shows a weak inhibition. Thus, the AID protein can considered to be a zinc-dependent cytidine deaminase, as is APOBEC-1.

(9) Strong expression of AID mRNA ~~expression~~ is seen in lymph nodes (mesenteric and amygdaline). Also, weak expression in spleen is seen.

(10) ~~mRNA~~ eExpression of AID protein mRNA is seen in a variety of lymphoid tissues (Peyer's patches, mesenteric lymph node, axillary lymph node, spleen, and bone marrow). Especially; notable expression is seen in peripheral

lymphoid organs, such as lymphatic nodes and Peyer's patches. ~~Contrariwise,~~ In contrast, expression in primary lymphoid organs is lower than the peripheral lymphoid organs.

(11) Expression of AID mRNA is at the lower limit of detection limit level without cytokines (IL-4, CD40L, TGF- $\beta$ ) stimulation in mouse B cell clone CH12F3-2, in which the cytokines stimulate class switch from IgM to IgA in the cells, ~~whereas the e~~ Expression is induced 3 hours after stimulation, and maximal expression is seen after 12 hours, with cytokine stimulation.

(12) AID mRNA expression in mouse B cell clone CH12F3-2 can be induced more strongly when stimulated with all three cytokines, IL-4, CD40L and TGF- $\beta$ , simultaneously, ~~rather than stimulated with any one of them alone.~~ Also, it can be considered that *de novo* protein synthesis is necessary for augmentation of AID mRNA expression, as the AID mRNA expression induction by cytokines in mouse B cell clone CH12F3-2 can be inhibited by cycloheximide, a protein synthesis inhibitor.

(13) ~~In the~~ in vitro tests, an augmentation of AID mRNA expression can be seen when normal mouse spleen B cells are stimulated with LPS alone, LPS+IL-4, or LPS+TGF- $\beta$ .

(14) ~~In the~~ in vivo tests, when normal mice are immunized with sheep red blood cells (SRBC), a significant augmentation of AID mRNA expression can be seen 5 days after immunization, in which SRBC are known to induce clonal expansion, germinal center formation, and class switch recombination and affinity maturation of immunoglobulin genes.

(15) The *in vivo* augmentation of AID mRNA expression by SRBC immunization is specifically seen in splenic CD19 positive B cells.

(16) AID mRNA expression in lymphoid organs is specifically seen in the germinal center, enriched with B cells activated by antigen stimulation.

(17) ~~The H~~ human AID gene locates is located at locus 12p13, close to the locus 12p13.1, where the APOBEC-1 gene is locates located.

[0017] According to the characteristics described above, the AID protein of the present invention can be considered to have a function of regulating various biological mechanisms

required for generation of antigen-specific immunoglobulins (specific antibodies), which eliminate non-self antigens (foreign antigen, self-reacting cells, etc.) that triggers various diseases. The mechanism for generation of immunoglobulin having high specificity to antigens includes germinal center functions such as activation of B cells, class switch recombination of immunoglobulin genes, somatic hypermutation, and affinity maturation. The AID protein of the present invention can be considered to be one of the enzymes that play an important role in the genetic editing occurring in germinal center B cells (e.g. class switch recombination and somatic mutation).

[0018] The dysfunction of the AID protein of the present invention can be the cause for the of humoral immunodeficiency since it induces failure of germinal center B cell function, such as antigen-specific B cell activation, class switch recombination, and somatic mutation. ~~Reversely, Conversely,~~ the hyperfunction of the AID protein may induce allergy disease or autoimmune disease since it can cause inappropriate B cell activation and needless class switch recombination and somatic mutation.

[0019] Therefore, regulation of the function of AID protein and the gene encoding it enables ~~preventing prevention and treating treatment of~~ various immunodeficiencies, autoimmune diseases, and allergies, which result from, for example, B cell dysfunctions (e.g., IgA deficiency, IgA nephropathy,  $\gamma$  globulinemia, hyper IgM syndrome, etc.) or class switch deficiency of immunoglobulin. Thus, the AID protein and the gene encoding the AID protein can be targets for the development of drugs for therapy of diseases mentioned above.

[0020] Examples of diseases whose onset prevention, symptom remission, therapy and/or symptomatic treatment effect is expected by regulating the function of the AID protein of the present invention or the gene encoding it include, for example, primary immunodeficiency syndrome with congenital disorder of immune system, mainly various immunodeficiencies considered to develop by B cell deficiency, decrease, or dysfunction (e.g., sex-linked agammaglobulinemia, sex-linked agammaglobulinemia with growth hormone deficiency, immunoglobulin deficiency with high IgM level, selective IgM deficiency, selective IgE deficiency, immunoglobulin heavy chain gene deletion,  $\kappa$  chain deficiency, IgA deficiency, IgG subclass selective deficiency, CVID (common variable immunodeficiency), infantile transient dysgammaglobulinemia, Rosen syndrome, severe combined immunodeficiency (sex-linked, autosomal recessive), ADA (adenosine deaminase) deficiency, PNP (purine

nucleoside phosphorylase) deficiency, MHC class II deficiency, reticular dysplasia, Wiskott-Aldrich syndrome, ataxia telangiectasia, DiGeorge syndrome, chromosomal aberration, familial Ig hypermetabolism, hyper IgE syndrome, Gitlin syndrome, Nezelof syndrome, Good syndrome, osteodystrophy, transcobalamin syndrome, secretory bead syndrome, etc.), various diseases with antibody production deficiency that are secondary immunodeficiency syndromes with a disorder of immune system caused by an acquired etiology (for example, AIDS, etc.), and/or various allergic diseases (e.g., bronchial asthma, atopic dermatitis, conjunctivitis, allergic rhinitis, allergic enteritis, drug-induced allergy, food allergy, allergic urticaria, glomerulonephritis, etc.).

[0021] Namely, ~~the~~ The AID proteins of the present invention, a fragment thereof, a DNA encoding the AID protein, a fragment thereof, and an antibody against the AID protein are useful as reagents for developing drugs for prevention and therapy of such diseases.

[0022] Also, the DNA itself is useful as an antisense drug regulating the function of the AID gene at a gene level and in ~~a use in~~ gene therapy. The protein or the fragments thereof (e.g. enzyme active site) ~~itself is themselves are~~ useful as ~~a~~ drugs.

[0023] Furthermore, a DNA comprising a ~~complementary~~ nucleotide sequence that is complementary to an arbitrary partial nucleotide sequence in the nucleotide sequence of genomic DNA encoding AID protein of the present invention (especially human AID protein) is useful as a primer DNA for polymerase chain reaction (PCR).

[0024] An arbitrary partial nucleotide sequence of genomic DNA encoding the AID protein (especially human AID protein) of the present invention can be amplified by PCR using the primer DNA pair. For example, in the case that mutation or deletion of the nucleotide sequence of genomic DNA (especially exon) encoding AID protein is presumed to cause a certain immunodeficiency or an allergy, mutations and deletions in the genomic DNA can be identified by amplifying an arbitrary partial nucleotide sequence of genomic DNA encoding the AID protein obtained from tissue or cells of immunodeficiency or allergy patients by PCR using a pair of ~~the~~ primer DNAs, by analyzing the presence and the size of PCR products; and the nucleotide sequence of the PCR products, and by comparing the nucleotide sequence with the corresponding nucleotide sequence in the genomic DNA encoding the AID protein derived from ~~the~~ a normal human. That is to say, this method is capable of not only, for example, elucidating relationships between immunodeficiency or allergy and AID



protein, but also, in the case where the AID protein is the cause of onset of a sort of disease (e.g. immunodeficiency and/or allergy), diagnosing the diseases by the methods mentioned above.

5 [0025] Furthermore, an antibody reactive to the AID protein of the present invention or a fragment thereof is extremely useful as an antibody drug by regulating functions of the AID protein.

[0026] Furthermore, the gene (DNA), protein, and antibody of the present invention are useful as reagents for ~~searching~~ searching for substrates (e.g. RNA, etc.) that ~~interacting~~ (binding) with the protein (enzyme) of the present invention, or other auxiliary proteins  
10 associated with the protein of the present invention, and for developing drugs targeting the substrates and auxiliary proteins.

[0027] Also, model animals can be generated by disrupting (inactivating) the AID gene based on the genetic information on the AID protein derived from mammals (e.g. mouse, etc.), which is one embodiment of the DNA of present invention. By analyzing the physical,  
15 biological, pathological, and genetic features of the model animal, it is possible to elucidate functions of the genes and the proteins of the present invention.

[0028] Furthermore, by introducing a normal human AID gene or mutant human AID gene (e.g. mutant human AID genes derived from immunodeficiency patients), which is one embodiment of the present invention, into the model animal whose endogenous gene has  
20 been disrupted, model animals having only normal or mutant human AID genes of the present invention can be generated. By administering drugs (compounds, antibodies, etc.) targeting the introduced human AID genes to the model animals, therapeutic effects of the drugs can be evaluated.

[0029] Furthermore, a method for identifying a substance that regulates production of the  
25 AID protein of the present invention or transcription of a gene encoding the AID protein into mRNA, or a substrate that inhibits the enzyme activity of the AID protein (e.g. cytidine deaminase activity) ~~are~~ is extremely useful as a means to develop drugs for therapy and prevention of various diseases (especially, immunodeficiency and/or allergy) in which the above-mentioned AID protein or AID gene is considered to be involved.

30 [0030] Thus, the present invention, for the first time, provides ~~below~~ the below-mentioned DNAs (cDNAs, genomic DNAs, and an arbitrary fragment thereof), proteins, expression

vectors, transformants, antibody pharmaceutical compositions, cells, the use of the DNA fragments as primer DNAs, and methods for screening.

(1) A DNA or a fragment thereof encoding a protein comprising the amino acid sequence of SEQ ID NO:2 or 8.

5 (2) The DNA or the fragment of (1), wherein the protein has a cytidine deaminase activity.

(3) A DNA or a fragment thereof comprising the nucleotide sequence of SEQ ID NO:1 or 7.

10 (4) A DNA or a fragment thereof comprising a nucleotide sequence of (a) or (b) below:

(a) a nucleotide sequence comprising the nucleotide residues 93 to 689 of SEQ ID NO:1 or

(b) a nucleotide sequence comprising the nucleotide residues 80 to 676 of SEQ ID NO:7.

15 (5) A DNA or a fragment thereof of (a) or (b) below:

(a) a DNA or a fragment thereof that hybridizes under stringent conditions with a DNA comprising the nucleotide sequence of SEQ ID NO:1 and that encodes a mammal-derived protein being homologous to a protein that comprises the amino acid sequence of SEQ ID NO:2 and having a cytidine deaminase activity or

20 (b) a DNA or a fragment thereof that hybridizes under stringent conditions with a DNA comprising the nucleotide sequence of SEQ ID NO:7 and that encodes a mammal-derived protein being homologous to a protein that comprises the amino acid sequence of SEQ ID NO:8 and having a cytidine deaminase activity.

25 (6) A protein or a fragment thereof comprising the amino acid sequence of SEQ ID NO:2 or 8.

(7) A protein or a fragment thereof comprising substantially the same amino acid sequence as that of SEQ ID NO:2 or 8 and having a cytidine deaminase activity.

(8) A proteins of (a) or (b) below.

30 (a) a mammal-derived protein that comprises an amino acid sequence encoded by a DNA hybridizing under stringent conditions with a DNA

comprising the nucleotide sequence of SEQ ID NO:1, that is homologous to a protein comprising the amino acid sequence of SEQ ID NO:2, and that has a cytidine deaminase activity, or

(b) a mammal-derived protein that comprises an amino acid sequence encoded by a DNA hybridizing under stringent conditions with a DNA comprising the nucleotide sequence of SEQ ID NO:7, that is homologous to a protein comprising the amino acid sequence of SEQ ID NO:8, and that has a cytidine deaminase activity.

(9) An expression vector comprising the DNA or the fragment of any one of (1) to (5).

(10) A transformant transformed with the expression vector of (9).

(11) An antibody or a portion thereof reactive to the protein of any one of (6) to (8) or to a fragment of the protein.

(12) The antibody or the portion thereof of (11), wherein the antibody is a monoclonal antibody.

(13) A pharmaceutical composition comprising the antibody or the portion thereof of (11) or (12), and a pharmaceutically acceptable carrier.

(14) A cell producing a monoclonal antibody reactive to the protein of any one of (6) to (8) or to a fragment of the protein.

(15) The cell of (14), wherein the cell is a hybridoma obtained by fusing, with a mammal-derived myeloma cell, a non-human mammal-derived B cell that produces a monoclonal antibody.

(16) The cell of (15), wherein the cell is a transgenic cell transformed by introducing, into a cell, either or both of a DNA encoding a heavy chain of the monoclonal antibody and a DNA encoding a light chain of the monoclonal antibody.

(17) A genomic DNA or a fragment thereof comprising a nucleotide sequence of any one of (a) to (c) below:

(a) SEQ ID NO:9,

(b) SEQ ID NO:10, or

(c) SEQ ID NO:35.

(18) A genomic DNA or a fragment thereof comprising a nucleotide sequence of any one of (a) to (e) below:

- (a) SEQ ID NO:11,
- (b) SEQ ID NO:12,
- (c) SEQ ID NO:13,
- (d) SEQ ID NO:14, or
- (e) SEQ ID NO:15.

(19) A DNA comprising a ~~complementary~~ nucleotide sequence complementary to an arbitrary partial nucleotide sequence of a nucleotide sequence of any one of (a) to (h) below:

- (a) SEQ ID NO:9,
- (b) SEQ ID NO:10,
- (c) SEQ ID NO:11,
- (d) SEQ ID NO:12,
- (e) SEQ ID NO:13,
- (f) SEQ ID NO:14,
- (g) SEQ ID NO:15, or
- (h) SEQ ID NO:25.

(20) The DNA of (19), wherein the DNA comprises a nucleotide sequence of any one of (a) to (q) below:

- (a) SEQ ID NO:18,
- (b) SEQ ID NO:19,
- (c) SEQ ID NO:20,
- (d) SEQ ID NO:21,
- (e) SEQ ID NO:22,
- (f) SEQ ID NO:23,
- (g) SEQ ID NO:24,
- (h) SEQ ID NO:25,
- (i) SEQ ID NO:26,
- (j) SEQ ID NO:27,
- (k) SEQ ID NO:28,

- 5
- (l) SEQ ID NO:29,
  - (m) SEQ ID NO:30,
  - (n) SEQ ID NO:31,
  - (o) SEQ ID NO:32,
  - (p) SEQ ID NO:33, or
  - (q) SEQ ID NO:34.

(21) Use of the DNA of (19) or (20) as a primer DNA in polymerase chain reaction.

10 (22) Use of a pair of DNAs of any one of (a) to (n) below as primer DNAs in polymerase chain reaction:

- (a) a DNA comprising the nucleotide sequence of SEQ ID NO:31 and a DNA comprising the nucleotide sequence of SEQ ID NO:32,
- (b) a DNA comprising the nucleotide sequence of SEQ ID NO:20 and a DNA comprising the nucleotide sequence of SEQ ID NO:22,
- 15 (c) a DNA comprising the nucleotide sequence of SEQ ID NO:21 and a DNA comprising the nucleotide sequence of SEQ ID NO:30,
- (d) a DNA comprising the nucleotide sequence of SEQ ID NO:24 and a DNA comprising the nucleotide sequence of SEQ ID NO:25,
- (e) a DNA comprising the nucleotide sequence of SEQ ID NO:23 and a DNA comprising the nucleotide sequence of SEQ ID NO:27,
- 20 (f) a DNA comprising the nucleotide sequence of SEQ ID NO:23 and a DNA comprising the nucleotide sequence of SEQ ID NO:28,
- (g) a DNA comprising the nucleotide sequence of SEQ ID NO:23 and a DNA comprising the nucleotide sequence of SEQ ID NO:29,
- 25 (h) a DNA comprising the nucleotide sequence of SEQ ID NO:26 and a DNA comprising the nucleotide sequence of SEQ ID NO:27,
- (i) a DNA comprising the nucleotide sequence of SEQ ID NO:26 and a DNA comprising the nucleotide sequence of SEQ ID NO:28,
- 30 (j) a DNA comprising the nucleotide sequence of SEQ ID NO:26 and a DNA comprising the nucleotide sequence of SEQ ID NO:29,

(k) a DNA comprising the nucleotide sequence of SEQ ID NO:34 and a DNA comprising the nucleotide sequence of SEQ ID NO:28,

(l) a DNA comprising the nucleotide sequence of SEQ ID NO:34 and a DNA comprising the nucleotide sequence of SEQ ID NO:29,

5 (m) a DNA comprising the nucleotide sequence of SEQ ID NO:33 and a DNA comprising the nucleotide sequence of SEQ ID NO:29, or,

(n) a DNA comprising the nucleotide sequence of SEQ ID NO:18 and a DNA comprising the nucleotide sequence of SEQ ID NO:19.

10 (23) A method for identifying a substance that regulates transcription of a gene encoding an AID protein comprising the amino acid sequence of SEQ ID NO:2 or 8 into mRNA, or production of the AID protein, the method comprising the steps of:

(a) culturing, separately in the presence and the absence of the substance, cells producing the AID protein and

15 (b) (i) comparing the level of the AID protein produced by the cells cultured in the presence of the substance with the level of the AID protein produced by the cells cultured in the absence of the substance or

(ii) comparing the level of the AID protein-encoding mRNA transcribed in the cells cultured in the presence of the substance with the level of the AID protein-encoding mRNA transcribed in the cells cultured in the absence or the substance.

20 (24) A method for identifying a substance that regulates transcription of a gene encoding an AID protein comprising the amino acid sequence of SEQ ID NO:2 or 8 into mRNA, or production of the AID protein, the method comprising the steps of:

25 (a) culturing, separately in the presence and the absence of the substance, cells producing the AID protein and a protein other than the AID protein, wherein transcription of a gene encoding the other protein into mRNA is dependent in the cells on the degree of a signal of transcription of the gene encoding the AID protein into mRNA and

30

(b) comparing the level of the other protein produced by the cells cultured in the presence of the substance with the level of the other protein produced by the cells cultured in the absence of the substance.

5 (25) The method of (23) or (24), wherein the cells are transgenic cells transformed with a gene encoding the protein.

(26) The method of (24), wherein the cells are transgenic cells transformed with a gene encoding the protein and a gene encoding the other protein.

(27) The method of (26), wherein the protein is a reporter protein.

10 (28) The method of (27), wherein comparison of the level of the other protein is comparison of the level of a signal generated by the reporter protein.

(29) The method of (27) or (28), wherein the reporter protein is luciferase.

(30) A method for identifying a substance that inhibits an enzyme activity of an AID protein comprising the amino acid sequence of SEQ ID NO:2 or 8, the method comprising the step of (a) or (b) below:

15 (a) culturing, separately in the presence and the absence of the substance, mammal-derived B cells or tissues comprising the B cells, and comparing enzyme activities of the AID protein in the B cells separately cultured or

(b) (i) administering the substance separately to an AID gene knockout mouse whose endogenous AID gene is inactivated so that transcription of the endogenous AID gene into mRNA is inhibited, and to a normal mouse and

20 (ii) comparing enzyme activities of the AID proteins in the B cells isolated from the respective mice.

(31) The method of (30), wherein the enzyme activity is a cytidine deaminase activity.

25 **[0031]** Hereafter, the present invention is explained in detail, by clarifying the terms used in the present invention and general methods for producing the proteins, DNAs, antibodies, and cells of the present invention.

**[0032]** The “protein or a fragment thereof” means a protein and a fragment thereof derived from a mammal such as human, bovine, sheep, pig, goat, rabbit, rat, hamster, guinea pig, mouse, and so on, preferably a protein or a fragment thereof derived from human, rabbit, rat,

or mouse, and particularly preferably, a protein ~~of~~or a fragment thereof derived from human or mouse.

[0033] As a particularly preferred embodiment, it means any protein or a fragment thereof below.

5                   (1)     A protein or a fragment thereof comprising the amino acid sequence of SEQ ID NO:2 or 8.

                  (2)     A protein or a fragment thereof comprising substantially the same amino acid sequence as that of SEQ ID NO:2 or 8 and having a cytidine deaminase activity.

10                  (3)     A mammal-derived protein that comprises an amino acid sequence encoded by a DNA hybridizing under stringent conditions with a DNA comprising the nucleotide sequence of SEQ ID NO:1, that is homologous to a protein comprising the amino acid sequence of SEQ ID NO:2, and that has a cytidine deaminase activity.

                  (4)     A mammal-derived protein that comprises an amino acid sequence  
15                  encoded by a DNA hybridizing under stringent conditions with a DNA comprising the nucleotide sequence of SEQ ID NO:7, that is homologous to a protein comprising the amino acid sequence of SEQ ID NO:8, and that has a cytidine deaminase activity.

[0034] Here, “having substantially the same amino acid sequence” means that a protein has an amino acid sequence where multiple amino acids, preferably 1 to 10 amino acids,  
20                  particularly preferably 1 to 5 amino acids, in the amino acid sequence shown in the references are substituted, deleted, and/or modified, and that a protein has an amino acid sequence where multiple amino acids, preferably 1 to 10 amino acids, particularly preferably 1 to 5 amino acids, are added to the amino acid sequence shown in the references.

[0035] The protein of the present invention includes monomer molecules, homodimers in  
25                  which one strand binds to another strand comprising an identical amino acid sequence, heterodimers in which one strand binds to another strand comprising a different amino acid sequence, and oligomers such as trimers or tetramers.

[0036] Also, ~~the~~a “fragment of a protein” means an arbitrary partial sequence (fragment) in  
30                  the amino acid sequence that the above-mentioned AID protein of the present invention comprises. For example, it includes an enzyme active site required for the AID protein to exert an enzyme activity represented by a cytidine deaminase activity, and an interaction site



required for the AID protein to bind or associate with substrates (e.g. mRNA, etc.) or various auxiliary proteins.

[0037] Alphabetical triplet or single letter codes used to represent amino acids in the present specification or figures mean amino acids as follows:

5 [0038] (Gly/G), glycine; (Ala/A), alanine; (Val/V), valine; (Leu/L), leucine; (Ile/I), isoleucine; (Ser/S), serine; (Thr/T), threonine; (Asp/D), aspartic acid; (Glu/E), glutamic acid; (Asn/N), asparagines; (Gln/Q) glutamine; (Lys/K), lysine; (Arg/R), arginine; (Cys/C), cysteine; (Met/M), methionine; (Phe/F), phenylalanine; (Tyr/Y), tyrosine; (Trp/W), tryptophan; (His/H), histidine; (Pro/P), proline.

10 [0039] The proteins and fragments of the present invention can be produced by properly using, in addition to genetic engineering technique mentioned below, methods well known in the art, such as chemical synthesis, cell culture method, and so on, or their modified methods.

[0040] Also, the AID protein of the present invention can be produced as a recombinant fusion protein with another protein (e.g. GST (Glutathione S-transferase), etc.). In this case,  
15 the fusion protein is advantageous in that it can be extremely easily purified by affinity chromatography employing adsorbent on which another ~~protein~~-molecule binding specifically to GST is immobilized. Moreover, since various antibodies reactive to GST are provided, the quantification of the fusion protein can be simply carried out by immunoassay (e.g. ELISA, etc.) using ~~the~~ antibodies against GST.

20 [0041] The DNA of the present invention is a DNA encoding a protein of the present invention and a fragment thereof, and it includes any nucleotide sequence encoding the protein of the present invention and includes both genomic DNAs and cDNAs. Also, the DNA includes any DNA composed of any codons as long as the codons encode identical amino acids.

25 [0042] Also, the DNA of the present invention includes a DNA encoding a mammalian AID protein, and, as a preferred embodiment, a DNA encoding a mouse AID protein or a human AID protein can be exemplified.

[0043] Examples of specific embodiments are as follows:

- 30 (1) A DNA encoding a protein comprising the amino acid sequence of SEQ ID NO:2 or 8.
- (2) The DNA of (1), wherein the protein has a cytidine deaminase activity.

- (3) A DNA comprising the nucleotide sequences of SEQ ID NO:1 or 7.
- (4) A DNA comprising ~~the nucleotides~~ residues 93 to 689 of SEQ ID NO:1.
- (5) A DNA comprising ~~the nucleotides~~ residues 80 to 676 of SEQ ID NO:7.
- 5 (6) A DNA that hybridizes under stringent conditions with a DNA comprising the nucleotide sequence of SEQ ID NO:1 and that encodes a mammal-derived protein being homologous to a protein that comprises the amino acid sequence of SEQ ID NO:2 and having a cytidine deaminase activity.
- (7) A DNA that hybridizes under stringent conditions with a DNA comprising the nucleotide sequence of SEQ ID NO:7 and that encodes a mammal-derived protein  
10 being homologous to a protein that comprises the amino acid sequence of SEQ ID NO:8 and having a cytidine deaminase activity.
- (8) A genomic DNA or a fragment thereof comprising a nucleotide sequence of any one of (a) to (c) below:
- 15 (a) SEQ ID NO:9,  
(b) SEQ ID NO:10, or  
(c) SEQ ID NO:35.
- (9) A genomic DNA or a fragment thereof comprising a nucleotide sequence of any one of (a) to (e) below:
- 20 (a) SEQ ID NO:11,  
(b) SEQ ID NO:12,  
(c) SEQ ID NO:13,  
(d) SEQ ID NO:14, or  
(e) SEQ ID NO:15.
- (10) A DNA comprising a complementary nucleotide sequence to an arbitrary  
25 partial sequence of a nucleotide sequence of any one of (a) to (h) below:
- 30 (a) SEQ ID NO:9,  
(b) SEQ ID NO:10,  
(c) SEQ ID NO:11,  
(d) SEQ ID NO:12,  
(e) SEQ ID NO:13,  
(f) SEQ ID NO:14,

(g) SEQ ID NO:15, or

(h) SEQ ID NO:35.

(11) A DNA comprising a nucleotide sequence of any one of (a) to (q) below:

(a) SEQ ID NO:18,

(b) SEQ ID NO:19,

(c) SEQ ID NO:20,

(d) SEQ ID NO:21,

(e) SEQ ID NO:22,

(f) SEQ ID NO:23,

(g) SEQ ID NO:24,

(h) SEQ ID NO:25,

(i) SEQ ID NO:26,

(j) SEQ ID NO:27,

(k) SEQ ID NO:28,

(l) SEQ ID NO:29,

(m) SEQ ID NO:30,

(n) SEQ ID NO:31,

(o) SEQ ID NO:32,

(p) SEQ ID NO:33, or,

(q) SEQ ID NO:34.

[0044] Furthermore, a DNA encoding a mutant protein or a fragment thereof obtained by substituting, deleting, and/or modifying multiple amino acids, preferably 1 to 10 amino acids, particularly preferably 1 to 5 amino acids, or by inserting multiple amino acids, preferably 1 to 10 amino acids, particularly preferably 1 to 5 amino acids in the amino acid sequence constituting the above-defined AID protein of the present invention or a fragment thereof is included in the DNA of the present invention.

[0045] The term “under stringent conditions” used herein means, for example, the following conditions. For example, in the case of carrying out hybridization using a probe with not less than 50 bases in 0.9% NaCl, target temperature of causing 50% dissociation ( $T_m$ ) can be calculated from the formula below, and the hybridization temperature can be set as the formula below.

$$T_m = 82.3^{\circ}\text{C} + 0.41 \times (\text{G+C})\% - 500/n - 0.61 \times (\text{formamide})\%$$

(n means the number of bases of the probe)

$$\text{Temperature} = T_m - 25^{\circ}\text{C}$$

[0046] Also, in the case of using a probe with not less than 100 bases (G+C = 40 to 50%),  
 5 the changes of  $T_m$  as (1) and (2) below can be used as the indicator.

- (1) Every 1% mismatch decreases  $T_m$  by approximately  $1^{\circ}\text{C}$ .
- (2) Every 1% formamide decreases  $T_m$  by  $0.6$  to  $0.7^{\circ}\text{C}$ .

[0047] Thus, the temperature condition in the case of combination of complete complementary strands can be set as below.

- 10 (A)  $65$  to  $75^{\circ}\text{C}$  (without formamide)
- (B)  $35$  to  $45^{\circ}\text{C}$  (with 50% formamide)

[0048] The temperature condition in the case of combination of incomplete complementary strands can be set as below.

- (A)  $45$  to  $55^{\circ}\text{C}$  (without formamide)
- 15 (B)  $35$  to  $42^{\circ}\text{C}$  (with 30% formamide)

[0049] In the case of using probes with not more than 23 bases, temperature can be  $37^{\circ}\text{C}$ , or the formula below can also be used as an indicator.

$$\text{Temperature} = 2^{\circ}\text{C} \times (\text{number of A+T}) + 4^{\circ}\text{C} \times (\text{number of C+G}) - 5^{\circ}\text{C}$$

[0050] The DNA of the present invention can be a DNA obtained by any method. For  
 20 example, the DNA includes complementary DNA (cDNA) prepared from mRNA, DNA prepared from genomic DNA, DNA prepared by chemical synthesis, DNA obtained by PCR amplification with RNA or DNA as a template, and DNA constructed by appropriately combining these methods.

[0051] As used herein, an "isolated nucleic acid" is a nucleic acid, the structure of which is  
 25 not identical to that of any naturally occurring nucleic acid or to that of any fragment of a naturally occurring genomic nucleic acid spanning more than three genes. The term therefore covers, for example, (a) a DNA which has the sequence of part of a naturally occurring genomic DNA molecule but is not flanked by both of the coding sequences that flank that part of the molecule in the genome of the organism in which it naturally occurs; (b)  
 30 a nucleic acid incorporated into a vector or into the genomic DNA of a prokaryote or eukaryote in a manner such that the resulting molecule is not identical to any naturally

occurring vector or genomic DNA; (c) a separate molecule such as a cDNA, a genomic fragment, a fragment produced by polymerase chain reaction (PCR), or a restriction fragment; and (d) a recombinant nucleotide sequence that is part of a hybrid gene, i.e., a gene encoding a fusion protein. Specifically excluded from this definition are nucleic acids

5 present in random, uncharacterized mixtures of different DNA molecules, transfected cells, or cell clones, e.g., as these occur in a DNA library such as a cDNA or genomic DNA library.

[0052] The term "substantially pure" as used herein in reference to a given polypeptide means that the polypeptide is substantially free from other biological macromolecules. For example, the substantially pure polypeptide is at least 75%, 80, 85, 95, or 99% pure by dry  
10 weight. Purity can be measured by any appropriate standard method known in the art, for example, by column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

[0053] The invention also includes a polypeptide, or fragment thereof, that differs from the corresponding sequence shown as SEQ ID NO:2 or 8. The differences are, preferably, differences or changes at a non-essential residue or a conservative substitution. In one

15 embodiment, the polypeptide includes an amino acid sequence at least about 60% identical to a sequence shown as SEQ ID NO:2 or 8, or a fragment thereof. Preferably, the polypeptide is at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or more identical to SEQ ID NO:2 or 8 and has at least one cytidine deaminase function or activity described herein.

Preferred polypeptide fragments of the invention are at least 10%, preferably at least 20%,  
20 30%, 40%, 50%, 60%, 70%, or more, of the length of the sequence shown as SEQ ID NO:2 or 8 and have at least one cytidine deaminase activity described herein. Or alternatively, the fragment can be merely an immunogenic fragment.

[0054] As used herein, "% identity" of two amino acid sequences, or of two nucleic acid sequences, is determined using the algorithm of Karlin and Altschul (PNAS USA 87:2264-  
25 2268, 1990), modified as in Karlin and Altschul, PNAS USA 90:5873-5877, 1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (J. Mol. Biol. 215:403-410, 1990). BLAST nucleotide searches are performed with the NBLAST program, score = 100, wordlength = 12. BLAST protein searches are performed with the XBLAST program, score = 50, wordlength = 3. To obtain gapped alignment for  
30 comparison purposes GappedBLAST is utilized as described in Altschul et al (Nucleic Acids Res. 25:3389-3402, 1997). When utilizing BLAST and GappedBLAST programs the default

parameters of the respective programs (e.g., XBLAST and NBLAST) are used to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention.

[0055] Accordingly, in one aspect, the invention provides an isolated or purified nucleic acid molecule that encodes a polypeptide described herein or a fragment thereof. Preferably, the isolated nucleic acid molecule includes a nucleotide sequence that is at least 60% identical to the nucleotide sequence shown in SEQ ID NO:1 or 7. More preferably, the isolated nucleic acid molecule is at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more, identical to the nucleotide sequence shown in SEQ ID NO:1 or 7. In the case of an isolated nucleic acid molecule which is longer than or equivalent in length to the reference sequence, e.g., SEQ ID NO:1 or 7, the comparison is made with the full length of the reference sequence. Where the isolated nucleic acid molecule is shorter than the reference sequence, e.g., shorter than SEQ ID NO:1 or 7, the comparison is made to a segment of the reference sequence of the same length (excluding any loop required by the homology calculation).

[0056] The DNA encoding the protein of the present invention can be prepared by the usual methods: cloning cDNA from mRNA encoding the protein of the present invention, isolating genomic DNA and splicing it, chemical synthesis, and so on.

[0057] (1) cDNA can be cloned from mRNA encoding the protein of the present invention by, for example, the method described below.

[0058] First, the mRNA encoding the protein of the present invention is prepared from the above-mentioned tissues or cells expressing and producing the protein of the present invention. mRNA can be prepared by isolating total RNA by a known method such as guanidine-thiocyanate method (Chirgwin et al., Biochemistry, 18:5294, 1979), hot phenol method, or AGPC method, and subjecting it to affinity chromatography using oligo-dT cellulose or poly-U Sepharose.

[0059] Then, with the mRNA obtained as a template, cDNA is synthesized, for example, by a well-known method using reverse transcriptase, such as the method of Okayama et al (Mol. Cell. Biol. 2:161 (1982); Mol. Cell. Biol. 3:280 (1983)) or the method of Hoffman et al. (Gene 25:263 (1983)), and converted into double-stranded cDNA. A cDNA library is

prepared by transforming *E. coli* with plasmid vectors, phage vectors, or cosmid vectors having this cDNA or by transfecting *E. coli* after *in vitro* packaging.

[0060] The plasmid vectors used in this invention are not limited as long as they are replicated and maintained in hosts. Any phage vector that can be replicated in hosts can also be used. Examples of usually used cloning vectors are pUC19,  $\lambda$ gt10,  $\lambda$ gt11, and so on.

When the vector is applied to immunological screening as mentioned below, a vector having a promoter that can express a gene encoding the desired protein in a host is preferably used.

[0061] cDNA can be inserted into a plasmid by, for example, the method of Maniatis et al.

(Molecular Cloning, A Laboratory Manual, second edition, Cold Spring Harbor Laboratory,

p.1.53, 1989). cDNA can be inserted into a phage vector by, for example, the method of Hyunh et al. (DNA cloning, a practical approach, 1, p.49 (1985)). These methods can be simply performed by using a commercially available cloning kit (for example, a product from Takara Shuzo). The recombinant plasmid or phage vector thus obtained is introduced into an appropriate host cell such as a prokaryote (for example, *E. coli*: HB101, DH5a, MC1061/P3, etc).

[0062] Examples of a method for introducing a plasmid into a host are, calcium chloride method, calcium chloride/rubidium chloride method and electroporation method, described in Molecular Cloning, A Laboratory Manual (second edition, Cold Spring Harbor Laboratory, p.1.74 (1989)). Phage vectors can be introduced into host cells by, for example, a method in which the phage DNAs are introduced into grown hosts after *in vitro* packaging. *In vitro* packaging can be easily performed with a commercially available *in vitro* packaging kit (for example, a product from Stratagene or Amersham).

[0063] The identification of cDNA encoding protein, its expression being augmented depending on the stimulation of cytokines like AID protein of the present invention, can be carried out by for example suppression subtractive hybridization (SSH)(Proc. Natl. Acad. Sci. USA, 93:6025-6030, 1996; Anal. Biochem., 240:90-97, 1996) taking advantage of suppressive PCR effect (Nucleic Acids Res., 23:1087-1088, 1995), using two cDNA libraries, namely, a cDNA library constructed from mRNA derived from stimulated cells (tester cDNA library) and ~~that~~ one constructed from mRNA derived from unstimulated cells (driver cDNA library).

[0064] The preparation of cDNA libraries required for subtraction cloning can be performed by using a commercially available kit, for example, PCR-Select Subtraction Kit (CLONTECH, cat: K1804-1). The experiment can be performed according to the ~~document of procedure~~ instructions accompanying ~~in~~ the kit.

5 [0065] An example of a practical experimental procedure is listed below, briefly.

[0066] PolyA<sup>+</sup> RNA is prepared from cells with or without stimulation with appropriate stimulant as previously reported ~~method~~ (Nucleic Acids Res., 26:911-918, 1998). Next, cDNA is prepared, using reverse transcriptase, from each polyA<sup>+</sup> RNA samples, as is the commonly used method. cDNA prepared from stimulated cells is used as tester cDNA and  
10 that prepared from unstimulated cells as driver cDNA.

[0067] According to the previous report mentioned above and experimental manuals accompanying ~~with the~~ kit, driver cDNA is added to tester cDNA to perform subtraction. The efficiency of subtraction is monitored by adding small amount of exogenous DNA as a control. After subtraction, the exogenous DNA is concentrated.

15 [0068] The subtracted cDNA is cloned into an appropriate plasmid expression vector to construct a plasmid library ~~by by~~ a commonly used method.

[0069] Similar to the previously reported method, many colonies are screened by differential hybridization method (Nucleic Acids Res., 26:911-918, 1998; RINSYO-MEN-EKI, 29:451-459, 1997). Here, as the hybridization probes, tester cDNA and driver cDNA mentioned  
20 above labeled with radioisotope can be used. Clones containing the objective DNA ~~and that~~ or containing exogenous DNA can be distinguished by hybridizing the exogenous DNA with replicant filters.

[0070] Objective cDNA or its fragment can be obtained by selecting clones giving strong signals against radiolabeled tester cDNA probe rather than radiolabeled driver cDNA probe.

25 [0071] Also, cDNA encoding the protein of the present invention can be accomplished by other general cDNA screening methods.

[0072] For instance, cDNA or ~~its~~ a fragment encoding the protein of the present invention cloned by subtraction cloning method mentioned above, or chemically synthesized oligonucleotides corresponding to an amino acid sequence of the protein of the present  
30 invention, are labeled with <sup>32</sup>P to make probes, then by well-known colony hybridization methods (Crunstein et al., Proc. Natl. Acid. Sci. USA, 72:3961, 1975) or plaque hybridization



methods (Molecular Cloning, A Laboratory Manual, second edition, Cold Spring Harbor Laboratory, p.2.108, 1989), commercial or originally prepared cDNA libraries can be screened. Furthermore, a method to amplify DNA including cDNA encoding the protein of the present invention by PCR, by constructing a pair of PCR primers based on cDNA or its  
5 fragment encoding the protein of the present invention isolated by the subtraction cloning mentioned above, can be listed.

[0073] When a cDNA library prepared using a cDNA expression vector is used, the desired clone can be screened by the antigen-antibody reaction using an antibody against the desired protein. A screening method using PCR ~~method~~ methodology is preferably used when many  
10 clones are subjected to screening.

[0074] The nucleotide sequence of the DNA thus obtained can be determined by the Maxam-Gilbert method (Maxam et al., Proc. Natl. Acad. Sci. USA, 74:560 (1977)) or the dideoxynucleotide synthetic chain termination method using phage M13 (Sanger et al., Proc. Natl. Acad. Sci. USA, 74:5463-5467 (1977)). The nucleotide sequence can be easily  
15 determined using a commercial DNA sequencer.

[0075] The whole or a part of the gene encoding the protein of the present invention can be obtained by excising the clone obtained as mentioned above with restriction enzymes and so on.

[0076] (2) Also, the DNA encoding the protein of the present invention can be isolated from ~~the~~ genomic DNA derived from the cells expressing the protein of the present invention as mentioned above by the following methods.

[0077] Such cells are solubilized preferably by SDS or proteinase K, and the DNAs are deproteinized by repeating phenol extraction. RNAs are digested preferably with  
25 ribonuclease. The DNAs obtained are partially digested with appropriate restriction enzymes, and the DNA fragments obtained are amplified with appropriate phage or cosmid to generate a library. Then, clones having the desired sequence are detected, for example, by using radioactively labeled DNA probes, and the whole or a portion of the gene encoding the protein of the present invention is obtained from the clones by excision with restriction  
30 enzymes, etc.

[0078] For example, cDNA encoding a human-derived protein can be obtained by preparing a cosmid library into which human genomic DNAs (chromosomal DNAs) are introduced (“Laboratory Manual Human Genome Mapping,” M. Hori and Y. Nakamura, eds.,

Maruzen), screening the cosmid library to obtain positive clones containing DNA

5 corresponding to the coding region of the desired protein, and screening the above cDNA library using the coding region DNA excised from the positive clones as a probe.

[0079] Also, the present invention relates to any fragment of DNA (cDNA, genomic DNA, etc.) encoding an AID protein (especially a human AID protein) of the present invention described above. DNA with a ~~complementary~~-nucleotide sequence complementary to any

10 nucleotide sequence of cDNA or genomic DNA is useful as a primer DNA in polymerase chain reaction (PCR). By PCR using a pair of ~~the~~-primer DNAs, any partial nucleotide sequence of genomic DNA encoding AID protein (especially human AID protein) of the present invention can be amplified.

[0080] For instance, in the case that mutation or deletion of genomic DNA (especially exon) encoding the AID protein is presumed to cause a certain immunodeficiency or allergy, the existence of such ~~the~~-a mutation or deletion can be analyzed by PCR described below.

(1) Prepare a pair of primers comprising ~~complementary~~-nucleotide sequence complementary to any partial nucleotide sequence of genomic DNA encoding an AID protein of the present invention.

20 (2) Amplify the objective partial nucleotide sequence of the genomic DNA using the pair of primers, using genomic DNA encoding AID protein obtained from tissue or cells of immunodeficiency or allergy patients as templates.

(3) Analyze the existence of PCR products and the nucleotide sequence of the PCR products, and identify the mutation and deletion in the genomic DNA by comparing the nucleotide sequence and corresponding nucleotide sequence of genomic DNA encoding AID protein derived from a normal human.

25 [0081] Thus, the method described above can not only elucidate, for example, the relationship between immunodeficiency and/or allergy and AID protein, but also be used for the diagnosis of a certain kind of disease, in the case that AID protein is the cause of the disease.

30 [0082] Examples of the nucleotide sequence of the primer DNA are as follows:

(1) A DNA comprising a complementary nucleotide sequence to an arbitrary partial sequence of a nucleotide sequence of any one of (a) to (h) below:

- (a) SEQ ID NO:9,
- (b) SEQ ID NO:10,
- (c) SEQ ID NO:11,
- (d) SEQ ID NO:12,
- (e) SEQ ID NO:13,
- (f) SEQ ID NO:14,
- (g) SEQ ID NO:15, or
- (h) SEQ ID NO:35.

(2) A DNA comprising a nucleotide sequence of any one of (a) to (q) below:

- (a) SEQ ID NO:18,
- (b) SEQ ID NO:19,
- (c) SEQ ID NO:20,
- (d) SEQ ID NO:21,
- (e) SEQ ID NO:22,
- (f) SEQ ID NO:23,
- (g) SEQ ID NO:24,
- (h) SEQ ID NO:25,
- (i) SEQ ID NO:26,
- (j) SEQ ID NO:27,
- (k) SEQ ID NO:28,
- (l) SEQ ID NO:29,
- (m) SEQ ID NO:30,
- (n) SEQ ID NO:31,
- (o) SEQ ID NO:32,
- (p) SEQ ID NO:33, or,
- (q) SEQ ID NO:34.

[0083] Also, the present invention relates to the use of the above-mentioned DNA fragment as a primer DNA in polymerase chain reaction.

[0084] Examples of the combination of primer DNAs for PCR in diagnosis accomplished by PCR gene amplification and by analyzing it are as follows:

- (1) a DNA comprising the nucleotide sequence of SEQ ID NO:31 and a DNA comprising the nucleotide sequence of SEQ ID NO:32,
- 5 (2) a DNA comprising the nucleotide sequence of SEQ ID NO:20 and a DNA comprising the nucleotide sequence of SEQ ID NO:22,
- (3) a DNA comprising the nucleotide sequence of SEQ ID NO:21 and a DNA comprising the nucleotide sequence of SEQ ID NO:30,
- (4) a DNA comprising the nucleotide sequence of SEQ ID NO:24 and a  
10 DNA comprising the nucleotide sequence of SEQ ID NO:25,
- (5) a DNA comprising the nucleotide sequence of SEQ ID NO:23 and a DNA comprising the nucleotide sequence of SEQ ID NO:27,
- (6) a DNA comprising the nucleotide sequence of SEQ ID NO:23 and a DNA comprising the nucleotide sequence of SEQ ID NO:28,
- 15 (7) a DNA comprising the nucleotide sequence of SEQ ID NO:23 and a DNA comprising the nucleotide sequence of SEQ ID NO:29,
- (8) a DNA comprising the nucleotide sequence of SEQ ID NO:26 and a DNA comprising the nucleotide sequence of SEQ ID NO:27,
- (9) a DNA comprising the nucleotide sequence of SEQ ID NO:26 and a  
20 DNA comprising the nucleotide sequence of SEQ ID NO:28,
- (10) a DNA comprising the nucleotide sequence of SEQ ID NO:26 and a DNA comprising the nucleotide sequence of SEQ ID NO:29,
- (11) a DNA comprising the nucleotide sequence of SEQ ID NO:34 and a DNA comprising the nucleotide sequence of SEQ ID NO:28,
- 25 (12) a DNA comprising the nucleotide sequence of SEQ ID NO:34 and a DNA comprising the nucleotide sequence of SEQ ID NO:29,
- (13) a DNA comprising the nucleotide sequence of SEQ ID NO:33 and a DNA comprising the nucleotide sequence of SEQ ID NO:29, or,
- (14) a DNA comprising the nucleotide sequence of SEQ ID NO:18 and a  
30 DNA comprising the nucleotide sequence of SEQ ID NO:19.

[0085] Moreover, the present invention also relates to a recombinant vector comprising the DNA encoding the protein of the present invention. As a recombinant vector of the present invention, any vector can be used as long as it is capable of retaining replication or self-multiplication in each host cell of prokaryotic and/or eukaryotic cells, including plasmid  
5 vectors and phage vectors.

[0086] The recombinant vector can easily be prepared by ligating the DNA encoding a protein of the present invention with a vector for recombination available in the art (plasmid DNA and bacteriophage DNA) by the usual method.

[0087] Specific examples of the vectors used for recombination ~~used~~ are *E. coli*-derived  
10 plasmids such as pBR322, pBR325, pUC12, pUC13, and pUC19, yeast-derived plasmids such as pSH19 and pSH15, and *Bacillus subtilis*-derived plasmids such as pUB110, pTP5, and pC194. Examples of phages are a bacteriophage such as  $\lambda$  phage, and an animal or insect virus (pVL1393, Invitrogen) such as a retrovirus, vaccinia virus, and nuclear polyhedrosis virus.

[0088] An expression vector is useful for expressing the DNA encoding the protein of the  
15 present invention and for producing the protein of the present invention. The expression vector is not limited as long as it expresses the gene encoding the protein of the present invention in various prokaryotic and/or eukaryotic host cells and produces this protein. Examples thereof are pMAL C2, pEF-BOS (Nucleic Acids Res. 18:5322 (1990) and so on),  
20 pME18S (Experimental Medicine: SUPPLEMENT, "Handbook of Genetic Engineering" (1992) and so on), etc.

[0089] Also, the protein of the present invention can be produced as a fusion protein with other proteins. It can be prepared as a fusion protein, for example, with GST (Glutathione S-transferase) by subcloning a cDNA encoding the protein of the present invention, for  
25 example, into plasmid pGEX4T1 (Pharmacia), by transforming *E. coli* DH5 $\alpha$ , and by culturing the transformant.

[0090] When bacteria, particularly *E. coli*, are used as host cells, an expression vector generally comprises, at least, a promoter/operator region, an initiation codon, the DNA encoding the protein of the present invention, termination codon, terminator region, and  
30 replicon.

[0091] When yeast, animal cells, or insect cells are used as hosts, an expression vector is preferably comprising, at least, a promoter, an initiation codon, the DNA encoding the protein of the present invention, and a termination codon. It may also comprise the DNA encoding a signal peptide, enhancer sequence, 5'- and 3'-untranslated region of the gene  
5 encoding the protein of the present invention, splicing junctions, polyadenylation site, selectable marker region, and a replicon. The expression vector may also contain, if required, a gene for gene amplification (marker) that is usually used.

[0092] A promoter/operator region to express the protein of the present invention in bacteria comprises a promoter, an operator, and a Shine-Dalgarno (SD) sequence (for example,  
10 AAGG). For example, when the host is *Escherichia*, it preferably comprises Trp promoter, lac promoter, recA promoter,  $\lambda$ PL promoter, lpp promoter, tac promoter, or the like.

Examples of a promoter to express the protein of the present invention in yeast are PH05 promoter, PGK promoter, GAP promoter, ADH promoter, and so on. When the host is *Bacillus*, examples thereof are SL01 promoter, SP02 promoter, penP promoter, and so on.

15 When the host is a eukaryotic cell such as a mammalian cell, examples thereof are SV40-derived promoter, retrovirus promoter, heat shock promoter, and so on, and preferably an SV-40 and/or retrovirus-derived one. As a matter of course, the promoter is not limited to the above examples. In addition, using an enhancer is effective for expression.

[0093] A preferable initiation codon is, for example, a methionine codon (ATG).

20 [0094] A commonly used termination codon (for example, TAG, TAA, TGA) is exemplified as a termination codon.

[0095] Usually, ~~used~~-natural or synthetic terminators are used as a terminator region.

[0096] A replicon means a DNA capable of replicating the whole DNA sequence in host cells, and includes a natural plasmid, an artificially modified plasmid (DNA fragment  
25 prepared from a natural plasmid), a synthetic plasmid, and so on. Examples of preferable plasmids are pBR322 or its artificial derivatives (DNA fragment obtained by treating pBR322 with appropriate restriction enzymes) for *E. coli*, yeast 2  $\mu$  plasmid or yeast chromosomal DNA for yeast, and pRSVneo ATCC 37198, pSV2dhfr ATCC 37145, pdBPV-MMTneo ATCC 37224, pSV2neo ATCC 37149, and such for mammalian cells.

30 [0097] An enhancer sequence, polyadenylation site, and splicing junction that are usually used in the art, such as those derived from SV40, can also be used.

[0098] A selectable marker usually employed can be used according to the usual method. Examples thereof are resistance genes for antibiotics, such as tetracycline, ampicillin, or kanamycin.

5 [0099] Examples of genes for gene amplification are dihydrofolate reductase (DHFR) gene, thymidine kinase gene, neomycin resistance gene, glutamate synthase gene, adenosine deaminase gene, ornithine decarboxylase gene, hygromycin-B-phosphotransferase gene, aspartate transcarbamylase gene, etc.

[00100] The expression vector of the present invention can be prepared by continuously and circularly linking at least the above-mentioned promoter, initiation codon, 10 DNA encoding the protein of the present invention, termination codon, and terminator region, to an appropriate replicon. If desired, appropriate DNA fragments (for example, linkers, restriction sites, and so on), can be used by the usual method such as digestion with a restriction enzyme or ligation using T4 DNA ligase.

[00101] Transformants of the present invention can be prepared by introducing the 15 expression vector mentioned above into host cells.

[00102] Host cells used in the present invention are not limited as long as they are compatible with an expression vector mentioned above and can be transformed. Examples thereof are various cells such as wild-type cells or artificially established recombinant cells usually used in the technical field of the present invention (for example, bacteria 20 (*Escherichia* and *Bacillus*), yeast (*Saccharomyces*, *Pichia*, and such), animal cells, or insect cells).

[00103] *E. coli* or animal cells are preferably used. Specific examples are *E. coli* (DH5 $\alpha$ , TB1, HB101, and such), mouse-derived cells (COP, L, C127, Sp2/0, NS-1, NIH 3T3, and such), rat-derived cells (PC12, PC12h), hamster-derived cells (BHK, CHO, and such), 25 monkey-derived cells (COS1, COS3, COS7, CV1, Velo, and such), and human-derived cells (Hela, diploid fibroblast-derived cells, myeloma cells, and HepG2, and such).

[00104] An expression vector can be introduced (transformed (transfected)) into host cells by known methods.

[00105] Transformation can be performed, for example, according to the method of 30 Cohen et al. (Proc. Natl. Acad. Sci. USA, 69:2110 (1972)), the protoplast method (Mol. Gen. Genet., 168:111 (1979)), or the competent method (J. Mol. Biol., 56:209 (1971)) when the

hosts are bacteria (*E. coli*, *Bacillus subtilis*, and such), the method of Hinnen et al. (Proc. Natl. Acad. Sci. USA, 75:1927 (1978)), or the lithium method (J. Bacteriol., 153:163 (1983)) when the host is *Saccharomyces cerevisiae*, the method of Graham (Virology, 52:456 (1973)) when the hosts are animal cells, and the method of Summers et al. (Mol. Cell. Biol., 3:2156-2165 (1983)) when the hosts are insect cells.

[00106] The protein of the present invention can be produced by cultivating transformants (in the following, this term includes transfectants) comprising an expression vector prepared as mentioned above in nutrient media.

[00107] The nutrient media preferably comprises a carbon source, an inorganic nitrogen source, or an organic nitrogen source necessary for the growth of host cells (transformants). Examples of the carbon source are glucose, dextran, soluble starch, and sucrose, and examples of the inorganic or organic nitrogen source are ammonium salts, nitrates, amino acids, corn steep liquor, peptone, casein, ~~meat~~meat extract, soy bean cake, and potato extract. If desired, they may comprise other nutrients (for example, an inorganic salt (for example, calcium chloride, sodium dihydrogenphosphate, and magnesium chloride), vitamins, antibiotics (for example, tetracycline, neomycin, ampicillin, kanamycin, and so on).

[00108] Cultivation is performed by a method known in the art. Cultivation conditions such as temperature, pH of the media, and cultivation time are selected appropriately so that the protein of the present invention is produced in large quantities.

[00109] Specific media and cultivation conditions used depending on host cells are illustrated below, but are not limited thereto.

[00110] When the hosts are bacteria, actinomycetes, yeasts, or filamentous fungi, liquid media comprising the nutrient source mentioned above are appropriate. ~~The m~~Media with a pH of 5 to 8 are preferably used.

[00111] When the host is *E. coli*, examples of preferable media are LB media, M9 media (Miller et al. Exp. Mol. Genet., Cold Spring Harbor Laboratory, p.431 (1972)), and so on. Using these media, cultivation can be performed usually at 14 to 43°C for about 3 to 24 hours with aeration and stirring, if necessary.

[00112] When the host is *Bacillus*, cultivation can be performed usually at 30 to 40°C for about 16 to 96 hours with aeration and stirring, if necessary.



[00113] When the host is yeast, an example of medium media is Burkholder minimal medium media (Bostian, Proc. Natl. Acad. Sci. USA, 77:4505 (1980)). The pH of the medium media is preferably 5 to 8. Cultivation can be performed usually at 20 to 35°C for about 14 to 144 hours with aeration and stirring, if necessary.

5 [00114] When the host is an animal cell, examples of media are MEM media containing about 5 to 20% fetal bovine serum (Science, 122:501 (1952)), DMEM media (Virology, 8:396 (1959)), RPMI1640 medium media (J. Am. Med. Assoc., 199:519 (1967)), 199 medium media (Proc. Soc. Exp. Biol. Med., 73:1 (1950)), and so on. The pH of the medium media is preferably about 6 to 8. Cultivation can be performed usually at about 30  
10 to 40°C for about 15 to 72 hours with aeration and stirring, if necessary.

[00115] When the host is an insect cell, an example of media-medium is Grace's media medium containing fetal bovine serum (Proc. Natl. Acad. Sci. USA, 82:8404 (1985)). The pH thereof is preferably about 5 to 8. Cultivation can be performed usually at about 20 to 40°C for 15 to 100 hours with aeration and stirring, if necessary.

15 [00116] The protein of the present invention can be produced by cultivating transformants, especially mammalian cells, as mentioned above and allowing them to secrete the protein into the culture supernatant.

[00117] A culture filtrate (supernatant) is obtained by a method such as filtration or centrifugation of the obtained culture, and the protein of the present invention is purified and  
20 isolated from the culture filtrate by methods commonly used in order to purify and isolate a natural or synthetic protein.

[00118] Examples of the isolation and purification method are a method utilizing solubility, such as salting out and solvent precipitation method; a method utilizing the difference in molecular weight, such as dialysis, ultrafiltration, gel filtration, and sodium  
25 dodecyl sulfate-polyacrylamide gel electrophoresis; a method utilizing charges, such as ion exchange chromatography and hydroxylapatite chromatography; a method utilizing specific affinity, such as affinity column chromatography; a method utilizing the difference in hydrophobicity, such as reverse phase high performance liquid chromatography; and a method utilizing the difference in isoelectric point, such as isoelectric focusing.

30 [00119] When the protein of the present invention exists in the periplasm or cytoplasm of cultured transformants (for example, *E. coli*), first, the cells are harvested by a usual

method such as filtration or centrifugation and suspended in appropriate buffer. After the cell wall and/or cell membrane of the cells and such are disrupted by ~~the a~~ method such as lysis with sonication, lysozyme, and freeze-thawing, the membrane fraction comprising the protein of the present invention is obtained by ~~the a~~ method such as centrifugation or  
5 filtration. The membrane fraction is solubilized with a detergent such as Triton-X100 to obtain the crude extract. Finally, the protein is isolated and purified from the crude extract by ~~the a~~ usual method as illustrated above.

[00120] By using a DNA (cDNA or genomic DNA) encoding a human-derived AID protein included in the protein of the present invention, transgenic non-human mammals  
10 secreting the human AID protein in their body can be prepared. Namely, by integrating the human-derived DNA into an endogenous locus of non-human mammals (e.g. mouse), the human AID protein of the present invention encoded by the DNA is expressed and secreted in their body. The transgenic non-human mammals are included in the present invention.

[00121] The transgenic non-human mammals can be prepared according to the method  
15 usually used for producing a transgenic animal (for example, see "Newest Manual of Animal Cell Experiment,"; LIC press, Chapter 7, pp.361-408, (1990)).

[00122] Specifically, for example, a transgenic mouse can be produced as follows. Embryonic stem cells (ES cells) obtained from normal mouse blastocysts are transformed with an expression vector in which the gene encoding the human AID protein of the present  
20 invention and a marker gene (for example, neomycin resistance gene) have been inserted in an expressible manner. ES cells in which the gene encoding the human AID protein of the present invention has been integrated into the endogenous gene are screened by a usual method based on expression of the marker gene. Then, the ES cells screened are microinjected into a fertilized egg (blastocyst) obtained from another normal mouse (Proc.  
25 Natl. Acad. Sci. USA, 77:7380-7384 (1980); U.S. Pat. No. 4,873,191).

[00123] The blastocyst is transplanted into the uterus of another normal mouse as the foster mother. Then, founder mice are born from the foster mother. By mating the founder mice with normal mice, heterozygous transgenic mice are obtained. By mating the heterozygous transgenic mice with each other, homozygous transgenic mice are obtained  
30 according to Mendel's laws.

[00124] Also, a so-called “knockout mouse” can be generated based on the nucleotide sequence of DNA encoding mouse AID protein included in the present invention. The “knockout mouse” in the present invention means the mouse in which the endogenous gene encoding the mouse AID protein of the present invention is knocked-out (inactivated). For example, it can be generated by positive-negative selection method applying homologous recombination (U.S. Pat. Nos. 5,464,764; 5,487,992; 5,627,059; Proc. Natl. Acad. Sci. USA, 86:8932-8935, 1989; Nature, 342:435-438, 1989; etc.), and such knockout mice are one embodiment of the present invention.

[00125] The “antibody” in the present invention means a polyclonal antibody (antiserum) or a monoclonal antibody, and preferably a monoclonal antibody.

[00126] Specifically, it includes an antibody reactive to the above-mentioned protein of the present invention and a fragment thereof.

[00127] The “antibody” of the present invention also includes a natural antibody that can be prepared by immunizing mammals such as mice, rats, hamsters, guinea pigs, or rabbits with the protein of the present invention (including natural, recombinant, and chemically synthesized protein and cell), a fragment thereof, or a transformant highly expressing the protein of interest by recombinant technology mentioned above; a chimeric antibody and a humanized antibody (CDR-grafted antibody) that can be produced by recombinant technology; and a human monoclonal antibody that can be produced by using human antibody-producing transgenic animals.

[00128] The monoclonal antibody includes those having any one of the isotypes of IgG, IgM, IgA, IgD, or IgE. IgG or IgM is preferable.

[00129] The polyclonal antibody (antiserum) or monoclonal antibody of the present invention can be produced by known methods. Namely, mammals, preferably, mice, rats, hamsters, guinea pigs, rabbits, cats, dogs, pigs, goats, horses, or ~~bovine~~cows, or more preferably, mice, rats, hamsters, guinea pigs, or rabbits are immunized, for example, with an antigen mentioned above with Freund’s adjuvant, if necessary. The polyclonal antibody can be obtained from the serum obtained from the animal so immunized. The monoclonal antibodies are produced as follows. Hybridomas are produced by fusing the antibody-producing cells obtained from the animal so immunized and myeloma cells incapable of producing autoantibodies. Then the hybridomas are cloned, and clones producing the

monoclonal antibodies showing the specific affinity to the antigen used for immunizing the mammal are screened.

[00130] Specifically, the monoclonal antibody can be produced as follows.

Immunizations are done by injecting or implanting once or several times the above-

5 mentioned protein of the present invention, a fragment thereof, the cells that express the protein, and so on, as an immunogen, if necessary, with Freund's adjuvant, subcutaneously, intramuscularly, intravenously, through the footpad, or intraperitoneally into mice, rats, hamsters, guinea pigs, or rabbits, preferably mice, rats or hamsters (including transgenic animals generated so as to produce antibodies derived from another animal such as ~~the a~~  
10 ~~transgenic mouse producing human antibodies~~). Usually, immunizations are performed ~~once~~ one to four times every one to fourteen days after the first immunization. Antibody-producing cells are obtained from the mammal so immunized in about one to five days after the last immunization.

[00131] Hybridomas that secrete a monoclonal antibody can be prepared by the

15 method of Köhler and Milstein (Nature, 256:495-497 (1975)) and by its modified method.

Namely, hybridomas are prepared by fusing antibody-producing cells contained in a spleen, lymph node, bone marrow, or tonsil obtained from the non-human mammal immunized as mentioned above, preferably a spleen, with myeloma cells without autoantibody-producing ability, which are derived from, preferably, a mammal such as mice, rats, guinea pigs,

20 hamsters, rabbits, or humans, or more preferably, mice, rats, or humans.

[00132] For example, mouse-derived myeloma P3/X63-AG8.653 (653; ATCC No.

CRL1580), P3/NSI/1-Ag4-1 (NS-1), P3/X63-Ag8.U1 (P3U1), SP2/0-Ag14 (Sp2/0, Sp2),

PAI, F0, or BW5147; rat-derived myeloma 210RCY3-Ag.2.3.; or human-derived myeloma

U-266AR1, GM1500-6TG-A1-2, UC729-6, CEM-AGR, D1R11, or CEM-T15 can be used as

25 a myeloma used for the cell fusion.

[00133] Hybridoma clones producing monoclonal antibodies can be screened by

cultivating the hybridomas, for example, in microtiter plates and by measuring the reactivity of the culture supernatant in the well in which hybridoma growth is observed, to the

immunogen used for the immunization mentioned above, for example, by an enzyme

30 immunoassay such as RIA and ELISA.

[00134] The monoclonal antibodies can be produced from hybridomas by cultivating the hybridomas *in vitro* or *in vivo* such as in the ascites of mice, rats, guinea pigs, hamsters, or rabbits, preferably mice or rats, more preferably mice and isolating the antibodies from the resulting the-culture supernatant or ascites fluid of a mammal.

5 [00135] *In vitro* cultivation can be performed depending on the property of cells to be cultured, on the object of a test study, and on various culture, by using known nutrient media or any nutrient media derived from known basal media for growing, maintaining, and storing the hybridomas to produce monoclonal antibodies in the culture supernatant.

[00136] Examples of basal media are low calcium concentration media such as  
10 Ham'F12 medium, MCDB153 medium, or low calcium concentration MEM medium, and high calcium concentration media such as MCDB104 medium, MEM medium, D-MEM medium, RPMI1640 medium, ASF104 medium, or RD medium. The basal media can contain, for example, sera, hormones, cytokines, and/or various inorganic or organic substances depending on the objective.

15 [00137] Monoclonal antibodies can be isolated and purified from the culture supernatant or ascites mentioned above by saturated ammonium sulfate precipitation, euglobulin precipitation method, caproic acid method, caprylic acid method, ion exchange chromatography (DEAE or DE52), affinity chromatography using anti-immunoglobulin column or protein A column.

20 [00138] Furthermore, monoclonal antibodies can be obtained in a large quantity by cloning a gene encoding a monoclonal antibody from the hybridoma, generating transgenic bovines, goats, sheep, or pigs in which the gene encoding the antibody is integrated in its endogenous gene using transgenic animal generating technique, and recovering the monoclonal antibody derived from the antibody gene from milk of the transgenic animals  
25 (Nikkei Science, No.4, pp.78-84 (1997)).

[00139] The "chimeric antibody" of the present invention means a monoclonal antibody prepared by genetic engineering, and specifically, a chimeric monoclonal antibody, for example, mouse/human chimeric antibody, whose variable region is a mouse immunoglobulin-derived variable region and whose constant region is a human  
30 immunoglobulin-derived constant region.

[00140] The constant region derived from human immunoglobulin has the amino acid sequence inherent in each isotype such as IgG, IgM, IgA, IgD, IgE, etc. The constant region of the recombinant chimeric monoclonal antibody of the present invention can be that of human immunoglobulin belonging to any isotype. Preferably, it is the constant region of human IgG.

[00141] The chimeric monoclonal antibody of the present invention can be produced, for example, as follows. Needless to say, the production method is not limited thereto.

[00142] For example, mouse/human chimeric monoclonal antibody can be prepared, by referring to Experimental Medicine: SUPPLEMENT, 1.6, No.10 (1988); and Examined Published Japanese Patent Application (JP-B) No. Hei 3-73280. Namely, it can be prepared by ligating CH gene (C gene encoding the constant region of H chain) obtained from the DNA encoding human immunoglobulin to the downstream of active VH genes (rearranged VDJ gene encoding the variable region of H chain) obtained from the DNA encoding mouse monoclonal antibody isolated from the hybridoma producing the mouse monoclonal antibody, and by ligating the CL gene (C gene encoding the constant region of L chain) obtained from the DNA encoding human immunoglobulin to the downstream of active VL genes (rearranged VJ gene encoding the variable region of L chain) obtained from the DNA encoding the mouse monoclonal antibody isolated from the hybridoma, and operably inserting those into the same or different vectors in an expressible manner, followed by transformation of host cells with the expression vector, and cultivation of the transformants.

[00143] Specifically, DNAs are first extracted from mouse monoclonal antibody-producing hybridoma by the usual method, digested with appropriate restriction enzymes (for example, EcoRI and HindIII), electrophoresed (using, for example, 0.7% agarose gel), and analyzed by Southern blotting. After the electrophoresed gel is stained, for example, with ethidium bromide and photographed, the gel is given marker positions, washed twice with water, and soaked in 0.25 M HCl for 15 minutes. Then, the gel is soaked in 0.4 N NaOH solution for 10 minutes with gentle stirring. The DNAs are transferred to a filter for 4 hours following the usual method. The filter is recovered and washed twice with 2 x SSC. After the filter is sufficiently dried, it is baked at 75°C for 3 hours, treated with 0.1 x SSC/0.1% SDS at 65°C for 30 minutes, and then soaked in 3 x SSC/0.1% SDS. The filter obtained is treated with prehybridization solution in a plastic bag at 65°C for 3 to 4 hours.

[00144] Next,  $^{32}\text{P}$ -labeled probe DNA and hybridization solution are added to the bag and reacted at  $65^{\circ}\text{C}$  about 12 hours. After hybridization, the filter is washed under an appropriate salt concentration, reaction temperature, and time (for example,  $2 \times \text{SSC}/0.1\%$  SDS, room temperature, 10 minutes). The filter is put into a plastic bag with a little  $2 \times \text{SSC}$ ,  
5 and subjected to autoradiography after the bag is sealed.

[00145] Rearranged VDJ gene and VJ gene encoding H chain and L chain of mouse monoclonal antibody respectively are identified by Southern blotting mentioned above. The region comprising the identified DNA fragment is fractionated by sucrose density gradient centrifugation and inserted into a phage vector (for example, Charon 4A, Charon 28,  
10  $\lambda\text{EMBL3}$ ,  $\lambda\text{EMBL4}$ , etc.). *E. coli* (for example, LE392, NM539, etc.) are transformed with the phage vector to generate a genomic library. The genomic library is screened by plaque hybridization such as the Benton-Davis method (Science, 196:180-182 (1977)) using appropriate probes (H chain J gene, L chain ( $\kappa$ ) J gene, etc.) to obtain positive clones comprising rearranged VDJ gene or VJ gene respectively. By making the restriction map and  
15 determining the nucleotide sequence of the clones obtained, it is confirmed that genes comprising the desired, rearranged  $V_H$  (VDJ) gene or  $V_L$  (VJ) gene have been obtained.

[00146] Separately, human CH gene and human CL gene used for chimerization are isolated. For example, when a chimeric antibody with human IgG1 is produced,  $C\gamma_1$  gene is isolated as a  $C_H$  gene, and C $\kappa$  gene is also isolated as a  $C_L$  gene, are isolated. These genes  
20 can be isolated from human genomic library with mouse  $C\gamma_1$  gene and mouse C $\kappa$  gene, corresponding to human  $C\gamma_1$  gene and human C $\kappa$  gene, respectively, as probes, taking advantage of the high homology between the nucleotide sequences of mouse immunoglobulin gene and that of human immunoglobulin gene.

[00147] Specifically, DNA fragments comprising human C $\kappa$  gene and an enhancer  
25 region are isolated from human  $\lambda$  Charon 4A HaeIII-AluI genomic library (Cell, 15:1157-1174 (1978)), for example, using a 3 kb HindIII-BamHI fragment from clone Ig146 (Proc. Natl. Acad. Sci. USA, 75:4709-4713 (1978)) and a 6.8 kb EcoRI fragment from clone MEP10 (Proc. Natl. Acad. Sci. USA, 78:474-478 (1981)) as probes. In addition, for example, after human fetal hepatocyte DNA is digested with HindIII and fractionated by  
30 agarose gel electrophoresis, a 5.9 kb fragment is inserted into  $\lambda 788$  and then human  $C\gamma_1$  gene is isolated with the probes mentioned above.

[00148] Using a mouse V<sub>H</sub> gene, mouse V<sub>L</sub> gene, human C<sub>H</sub> gene, and human C<sub>L</sub> gene so obtained, and taking the promoter region and enhancer region into consideration, human C<sub>H</sub> gene is inserted downstream of mouse V<sub>H</sub> gene and human C<sub>L</sub> gene is inserted downstream of mouse V<sub>L</sub> gene in an expression vector such as pSV2gpt or pSV2neo with  
5 appropriate restriction enzymes and DNA ligase following the usual method. In this case, chimeric genes of mouse V<sub>H</sub> gene/human C<sub>H</sub> gene and mouse V<sub>L</sub> gene/human C<sub>L</sub> gene can be respectively inserted into ~~a~~the same or a different expression vector.

[00149] Chimeric gene-inserted expression vector(s) thus prepared are introduced into myeloma cells (e.g., P3X63 Ag8 653 cells or SP210 cells) that do not produce antibodies by  
10 the protoplast fusion method, DEAE-dextran method, calcium phosphate method, or electroporation method. The transformants are screened by cultivating them in a medium containing a drug corresponding to the drug resistance gene inserted into the expression vector and, then, cells producing the desired chimeric monoclonal antibodies are obtained.

[00150] Desired chimeric monoclonal antibodies are obtained from the culture  
15 supernatant of antibody-producing cells thus screened.

[00151] The “humanized antibody (CDR-grafted antibody)” of the present invention is a monoclonal antibody prepared by genetic engineering and specifically means a humanized monoclonal antibody wherein a portion or the whole of the complementarity determining regions of the hyper-variable region are derived from those of the hyper-variable region from  
20 mouse monoclonal antibody, the framework regions of the variable region are derived from those of the variable region from human immunoglobulin, and the constant region is derived from that from human-immunoglobulin.

[00152] The complementarity determining regions of the hyper-variable region exists in the hyper-variable region in the variable region of an antibody and means three regions  
25 which directly binds, in a complementary manner, to an antigen (complementarity-determining residues, CDR1, CDR2, and CDR3). The framework regions of the variable region mean four comparatively conserved regions intervening upstream, downstream or between the three complementarity-determining regions (framework region, FR1, FR2, FR3, and FR4).

[00153] In other words, a humanized monoclonal antibody means that in which the  
30 whole region except a portion or the whole region of the complementarity determining



regions of the hyper-variable region of a mouse monoclonal antibody has been replaced with their corresponding regions derived from human immunoglobulin.

[00154] The constant region derived from human immunoglobulin has the amino acid sequence inherent in each isotype such as IgG, IgM, IgA, IgD, and IgE. The constant region  
5 of a humanized monoclonal antibody ~~in~~of the present invention can be that from human immunoglobulin belonging to any isotype. Preferably, it is the constant region of human IgG. The framework regions of the constant region derived from human immunoglobulin are not particularly limited.

[00155] The humanized monoclonal antibody of the present invention can be  
10 produced, for example, as follows. Needless to say, the production method is not limited thereto.

[00156] For example, a recombinant humanized monoclonal antibody derived from mouse monoclonal antibody can be prepared by genetic engineering, referring to Published Japanese Translations of PCT International Publication No. Hei 4-506458 and Unexamined  
15 Published Japanese Patent Application (JP-A) No. Sho 62-296890. Namely, at least one mouse H chain CDR gene and at least one mouse L chain CDR gene corresponding to the mouse H chain CDR gene are isolated from hybridomas producing mouse monoclonal antibody, and human H chain gene encoding the whole region except human H chain CDR corresponding to mouse H chain CDR mentioned above and human L chain gene encoding  
20 the whole region except human L chain CDR corresponding to mouse L chain CDR mentioned above are isolated from human immunoglobulin genes.

[00157] The mouse H chain CDR gene(s) and the human H chain gene(s) so isolated are inserted, in an expressible manner, into an appropriate vector so that they can be expressed. Similarly, the mouse L chain CDR gene(s) and the human L chain gene(s) are  
25 inserted, in an expressible manner, into another appropriate vector so that they can be expressed. Alternatively, the mouse H chain CDR gene(s)/human H chain gene(s) and mouse L chain CDR gene(s)/human L chain gene(s) can be inserted, in an expressible manner, into the same expression vector so that they can be expressed. Host cells are transformed with the expression vector thus prepared to obtain transformants producing humanized monoclonal  
30 antibody. By cultivating the transformants, desired humanized monoclonal antibody is obtained from culture supernatant.

[00158] The “human antibody” used in the present invention is immunoglobulin in which the entire regions comprising the variable and constant region of the H chain, and the variable and constant region of the L chain constituting immunoglobulin are derived from the genes encoding human immunoglobulin.

5 [00159] The human antibody can be produced in the same way as the production method of polyclonal or monoclonal antibodies mentioned above by immunizing, with an antigen, a transgenic animal which for example, at least human immunoglobulin gene(s) have been integrated into the locus of a non-human mammal such as a mouse by the usual method.

[00160] For example, a transgenic mouse producing human antibodies is prepared by  
10 the methods described in already published literatures (Nature Genetics, 7:13-21 (1994); Nature Genetics, 15:146-156 (1997); JP-WA Hei 4-504365; WO94/25585; Nikkei Science, No.6, pp.40-50 (1995); WO94/25585; Nature, 368:856-859 (1994); JP-WA No. Hei 6-500233).

[00161] The “portion of an antibody” used in the present invention means a partial  
15 region of the antibody, and preferably the monoclonal antibody of the present invention as mentioned above, and specifically, means  $F(ab')_2$ ,  $Fab'$ ,  $Fab$ ,  $Fv$  (variable fragment of antibody),  $sFv$ ,  $dsFv$  (disulfide stabilized  $Fv$ ), or  $dAb$  (single domain antibody) (Exp. Opin. Ther. Patents, 6, No.5, pp.441-456 (1996)).

[00162] “ $F(ab')_2$ ” and “ $Fab'$ ” can be produced by treating immunoglobulin  
20 (monoclonal antibody) with a protease such as pepsin and papain, and means an antibody fragment generated by digesting immunoglobulin near the disulfide bonds existing between the hinge regions in each of the two H chains. For example, papain cleaves IgG upstream of the disulfide bonds existing between the hinge regions in each of the two H chains to generate two homologous antibody fragments in which an L chain composed of  $V_L$  (L chain variable region) and  $C_L$  (L chain constant region), and an H chain fragment composed of  $V_H$  (H chain variable region) and  $C_{H\gamma 1}$  ( $\gamma 1$  region in the constant region of H chain) are  
25 connected at their C terminal regions through a disulfide bond. Each of these two homologous antibody fragments is called  $Fab'$ . Pepsin also cleaves IgG downstream of the disulfide bonds existing between the hinge regions in each of the two H chains to generate an  
30 antibody fragment slightly larger than the fragment in which the two above-mentioned  $Fab'$  are connected at the hinge region. This antibody fragment is called  $F(ab')_2$ .

[00163] The “cell producing a monoclonal antibody reactive to a protein or a fragment thereof” of the present invention means any cell producing the above-described monoclonal antibody of the present invention.

[00164] More specifically, the following is included:

- 5 (1) B cells that are obtained by immunizing the non-human mammals with the above-mentioned protein of the present invention, a fragment thereof, or the cells producing the protein and that produce a monoclonal antibody reactive to the protein of the present invention or a fragment thereof.
- (2) The above-mentioned hybridomas (fused cell) prepared by fusing the thus-  
10 obtained B cells producing the antibody with myeloma cells derived from mammals.
- (3) Monoclonal antibody-producing transformants obtained by transforming cells  
other ~~cells~~ than the monoclonal antibody-producing B cells and hybridomas with  
genes encoding the monoclonal antibody isolated from the monoclonal antibody-  
producing B cells or hybridomas (either the heavy chain-encoding gene or the light  
15 chain-encoding gene, or both).

The monoclonal antibody-producing transformants of (3) mean recombinant cells producing a recombinant monoclonal antibody produced by B cells of (1) or hybridomas of (2). These antibody producing-transformants can be produced by the method as used for producing the above-described chimeric monoclonal antibody and  
20 humanized monoclonal antibody.

[00165] The “pharmaceutical composition” used herein means a pharmaceutical composition comprising of any of the protein, fragment thereof, antibody, or portion thereof defined hereinabove, and a pharmaceutically acceptable carrier.

[00166] The “pharmaceutically acceptable carrier” includes an excipient, a diluent, an  
25 expander, a disintegrating agent, a stabilizer, a preservative, a buffer, an emulsifier, an aromatic, a colorant, a sweetener, a viscosity-increasing agent, a flavor, a dissolving agent, or other additives. Using one or more of such carriers, a pharmaceutical composition can be formulated into tablets, pills, powders, granules, injections, solutions, capsules, troches, elixirs, suspensions, emulsions, or syrups. The pharmaceutical composition can be  
30 administered orally or parenterally. Other forms for parenteral administration include a

solution for external application, suppository for rectal administration, and pessary, prescribed by the usual method, which comprises one or more active ingredient.

5 [00167] The dosage can vary depending on the age, sex, weight, and symptoms of a patient, effect of treatment, administration route, period of treatment, or the kind of active ingredient (protein or antibody mentioned above) contained in the pharmaceutical composition. Usually, the pharmaceutical composition can be administered to an adult in a dose of 10  $\mu$ g to 1000 mg (or 10  $\mu$ g to 500 mg) per one administration. Depending on various conditions, the lower dosage may be sufficient in some cases, and a higher dosage may be necessary in other cases.

10 [00168] In particular, the injection can be produced by dissolving or suspending the antibody in a non-toxic, pharmaceutically acceptable carrier such as physiological saline or commercially available distilled water for injections by adjusting the concentration to 0.1  $\mu$ g antibody/ml carrier to 10 mg antibody/ml carrier. The injection thus produced can be administered to a human patient in need of treatment in a dose of 1  $\mu$ g to 100 mg/kg body weight, preferably 50  $\mu$ g to 50 mg/kg body weight, once or more times a day. Examples of administration routes are medically appropriate administration routes such as intravenous injection, subcutaneous injection, intradermal injection, intramuscular injection, or intraperitoneal injection, preferably intravenous injection.

20 [00169] The injection can also be prepared into a non-aqueous diluent (for example, propylene glycol, polyethylene glycol, vegetable oil such as olive oil, and alcohols such as ethanol), suspension, or emulsion.

[00170] The injection can be sterilized by filtration with a bacteria-non-penetrable filter, by mixing bactericide, or by irradiation. The injection can be prepared at the time of use. Namely, it is freeze-dried to make a sterile solid composition, and can be dissolved in sterile distilled water for injection or another solvent before use.

25 [00171] The pharmaceutical composition of the present invention is useful as a drug for preventing and treating, for example, primary immunodeficiency syndrome with congenital disorder of immune system, mainly immunodeficiency considered to develop by B lymphocyte deficiency, decrease, or dysfunction (e.g., sex-linked agammaglobulinemia, sex-linked agammaglobulinemia with growth hormone deficiency, immunoglobulin deficiency with high IgM level, selective IgM deficiency, selective IgE deficiency,

30

immunoglobulin heavy chain gene deletion,  $\kappa$  chain deficiency, IgA deficiency, IgG subclass selective deficiency, CVID (common variable immunodeficiency), infantile transient dysgammaglobulinemia, Rosen syndrome, severe combined immunodeficiency (sex-linked, autosomal recessive), ADA (adenosine deaminase) deficiency, PNP (purine nucleoside phosphorylase) deficiency, MHC class II deficiency, reticular dysplasia, Wiskott-Aldrich syndrome, ataxia telangiectasia, DiGeorge syndrome, chromosomal aberration, familial Ig hypermetabolism, hyper IgE syndrome, Gitlin syndrome, Nezelof syndrome, Good syndrome, osteodystrophy, transcobalamin syndrome, secretory bead syndrome, etc.), various diseases with antibody production deficiency that are secondary immunodeficiency syndrome with disorder of immune system caused by an acquired etiology (for example, AIDS, etc.), and/or various allergic diseases (e.g., bronchial asthma, atopic dermatitis, conjunctivitis, allergic rhinitis, allergic enteritis, drug-induced allergy, food allergy, allergic urticaria, glomerulonephritis, etc.), and for ~~reliving~~ relieving conditions due to various ~~immunodeficiency~~ immunodeficiencies associated with the diseases.

[00172] The DNA of the present invention described above, namely, "DNA comprising any partial nucleotide sequence of SEQ ID NO:7, from SEQ ID NO:9 to SEQ ID NO:15, or SEQ ID NO:35, those with partial chemical modification, DNA comprising complementary nucleotide sequences to the partial sequence, or those with partial chemical modification" are included.

[00173] Here, the "partial nucleotide sequence" means the partial nucleotide sequence comprising any number of bases at any region included in any nucleotide sequence listed in SEQ ID NO:7, from SEQ ID NO:9 to SEQ ID NO:15, or SEQ ID NO:35.

[00174] The DNA is useful as probes in DNA hybridization or RNA hybridization procedures. ~~In-For~~ For the purpose of using the DNA as a probe, continuous nucleotide sequences of over 20 bases, preferably continuous nucleotide sequences of over 50 bases, more preferably over 100 bases, much more preferably over 200 bases, especially preferably over 300 bases, can be ~~listed~~ used as the partial nucleotide sequences.

[00175] Also, the DNA described ~~above~~ above, as mentioned before, are useful as primers for PCR. ~~In-For~~ For the purpose of using the DNA as PCR primers, continuous partial nucleotide sequences of from 5 to 100 bases, preferably from 5 to 70 bases, more preferably

from 5 to 50 bases, much more preferably from 5 to 30 bases, can be ~~listed~~used as the partial nucleotide sequences.

[00176] Moreover, the DNA described above are useful as antisense drug. The DNA, by hybridizing to a DNA or an RNA encoding the AID protein of the present invention, can  
5 inhibit transcription of the DNA to mRNA or translation of the mRNA into the protein.

[00177] ~~In~~For the purpose of using above-mentioned DNA to antisense drug, the partial nucleotide sequence consists of 5 to 100 consecutive nucleotides, preferably 5 to 70 consecutive nucleotides, more preferably 5 to 50 consecutive nucleotides, and still more preferably 5 to 30 consecutive nucleotides.

10 [00178] When the DNA is used as an antisense DNA pharmaceutical, the DNA sequence can be modified chemically in part for extending the half-life (stability) of the blood concentration of the DNA administered to patients, for increasing the intracellular-membrane permeability of the DNA, or for increasing the degradation resistance or the absorption of the orally administered DNA in the digestive organs. The chemical  
15 modification includes, for example, the modification of the phosphate bonds, the riboses, the nucleotide bases, the sugar moiety, the 3' end and/or the 5' end in the structure of the oligonucleotide DNA.

[00179] The modification of phosphate bonds includes, for example, the conversion of one or more of the bonds to phosphodiester bonds (D-oligo), phosphorothioate bonds,  
20 phosphorodithioate bonds (S-oligo), methyl phosphonate (MP-oligo), phosphoroamidate bonds, non-phosphate bonds or methyl phosphonothioate bonds, or combinations thereof. The modification of the ribose includes, for example, the conversion to 2'-fluororibose or 2'-O-methylribose. The modification of the nucleotide base includes, for example, the conversion to 5-propynyluracil or 2-aminoadenine.

25 [00180] Also, another ~~one~~embodiment of the present invention relates to "methods of identifying substances regulating the production of the AID protein of the present invention or the transcription of the gene encoding AID protein to mRNA." The method of the present invention is namely "the method of screening of drugs capable of regulating functions of AID protein or AID gene".

30 [00181] As the cells ~~used~~ in the method of the present invention, any cells, as long as capable of producing AID protein of the present invention, can be used. For instance, native

cells (preferably of mouse or human), transgenic cells transformed with a gene encoding an AID protein of the present invention, cells introduced with RNA encoding an AID protein of the present invention, etc., can be ~~listed~~used.

5 [00182] As the host cells ~~used~~ for preparing the transgenic cells, various cells, mentioned in the part explaining in detail ~~on~~ the methods of expressing the protein of the present invention using the DNA of the protein described above, can be used.

[00183] For instance, various cells such as naturally established cells or artificially established transgenic cells (e.g. bacteria (*Escherichia*, *Bacillus*), yeast (*Saccharomyces*, *Pichia*), animal cells and insect cells) can be exemplified.

10 [00184] Preferably, animal cells, namely, cells derived from mouse (COP, L, C127, Sp2/0, NS-1, or NIH3T3, etc.), cells derived from rat (PC12, PC12h, etc.), cells derived from hamster (BHK, and CHO, etc.), cells derived from monkey (COS1, COS3, COS7, CV1, and Velo, etc.), and cells derived from human (Hela, cells derived from diploid fibroblast, HEK293 cells, myeloma cells, and Namalwa, etc.) can be exemplified.

15 [00185] The “substance” in the present invention means natural substance existing in the nature and any substance prepared artificially. The substances can be grouped into “peptidic substance” and “non-peptidic substance.”

[00186] As the “non-peptidic substance,” “DNA comprising partial nucleotide sequence, or chemically modified DNA derived from it” that are useful as antisense drugs as  
20 described above, “antisense RNA” with similar structural and pharmacological property to the antisense DNA, or any chemically synthesized “compounds” can be exemplified. The “compounds” herein means ~~compounds excepting DNA, RNA, and peptidic substances.~~  
~~Namely, compounds with molecular weight of smaller than from 100 to approximately 1000,~~  
~~preferably compounds with molecular weight of from 100 to 800, more preferably molecular~~  
25 ~~weight of from 100 to 600, can be exemplified.~~ Examples of “compounds” are compounds other than DNA, RNA, and the above-mentioned peptidic substances, which have a molecular weight from approximately 100 to approximately 1000, preferably from approximately 100 to approximately 800, more preferably from approximately 100 to approximately 600.

30 [00187] As the “peptidic substance,” antibodies already described above in detail (preferably monoclonal antibodies, more preferably recombinant antibodies or human

monoclonal antibodies), oligopeptides, or chemically modified substance derived from them can be exemplified. Examples of an oligopeptide are a peptide comprising 5 to 30 amino acids, preferably 5 to 20 amino acids. The chemical modification can be designed depending on various purposes, for example, ~~the~~for increased half-life in blood in the case of  
5 administering *in vivo*, or ~~the~~for increased tolerance against ~~the~~ degradation or increased absorption in digestive tract ~~at the~~after oral administration.

[00188] Methods described in from (24) to (28) above includes so-called reporter gene assays, as one of the method of the present invention.

[00189] As the “reporter protein<sub>1</sub>”, luciferase derived from firefly or sea pansy, or GFP  
10 derived from ~~jerry fish~~jellyfish, are preferred.

[00190] As the “reporter gene assay<sub>1</sub>”, methods described below are representative.

[00191] Transgenic cells are generated by transforming cells commonly used in the production of recombinant proteins with expression vector, in which a gene encoding the target protein and a gene encoding the reporter protein are inserted ~~intoto~~ the vector so that  
15 the transcription of the gene encoding the reporter protein to mRNA ~~occurs dependently~~  
~~depends on~~is induced by the signal of the transcription of the gene of target protein to mRNA. The test substances (described above) are applied to the obtained transformant cells. Analysis that whether the compound affects the expression of transporter molecule can be accomplished by measuring the level of the target protein by indirect ~~measuring~~  
20 measurement of the amount of fluorescence emitted by the reporter protein expressed in parallel with the target protein (for reference, see U.S. Pat. No. 5,436,128 and U.S. Pat. No. 5,401,629).

[00192] ~~Also,~~The identification of the compounds using the present assay can be accomplished by manual operation, but it can also be readily and simply done automatically  
25 by ~~using~~ so-called High-Throughput Screening using robots (SOSHIKI BAIYO KOUGAKU, 23:521-524; U.S. Pat. No. 5,670,113).

[00193] ~~The~~The terms “cells” and “substances” used in the methods described above ~~contain~~have the same meaning as defined above.

[00194] The substances identified by the methods of the present invention are very  
30 useful as drugs for the therapy of various diseases considered to be caused by the hyperfunction or deficiency of the AID protein of the present invention or by the deficiency



or mutation of the AID gene, or for remission of various symptoms ~~supervene~~ associated with the diseases.

### DESCRIPTION OF DRAWINGS

FIG. 1 is ~~a-is-the~~ a pair of photographs which shows the production ~~state-of~~ of DNA including ~~an Sa sequence~~ an Sa sequence looped out ~~with-the~~ by class switch recombination in mouse B cell clone CH12F3-2 cultured under ~~the~~ various conditions.

FIG. 1(a) shows the ~~electrophoretic-electrophoresis~~ state of DNA including ~~an Sa~~ an Sa sequence looped out ~~with-by~~ by class switch recombination, amplified by PCR using DNA derived from mouse B cell clone CH12F3-2 cultured under ~~the~~ various conditions.

10 Lanes 1 and 6 show the ~~electrophoretic-electrophoresis~~ state of marker DNAs. Lane 2 shows the ~~electrophoretic-electrophoresis~~ state of PCR products produced using DNA from cells cultured in ~~the~~ conditions excluding IL-4, CD40L, TGF  $\beta$  ~~or-and~~ and cycloheximide; as a template. Lane 3 shows the ~~electrophoretic-electrophoresis~~ state of DNA-PCR products produced using DNA from cells cultured in the presence of cycloheximide only, as a  
15 template. Lane 4 shows the ~~electrophoresistie~~ state of PCR products produced using DNA from cells cultured in the presence of IL-4, CD40L and TGF  $\beta$ ; as a template. Lane 5 shows the ~~electrophoretic-electrophoresis~~ state of PCR products produced using DNA from cells cultured in the presence of IL-4, CD40L, TGF  $\beta$ , and cycloheximide; as a template.

FIG. 1(b) shows the result of Southern hybridization ~~for-of~~ for DNA including an Sa  
20 sequence looped out ~~with-by~~ by class switch recombination, amplified by PCR using DNA derived from mouse B cell clone CH12F3-2 cultured under ~~the~~ various conditions.

Lane 1 shows the result of hybridization ~~against-of~~ against PCR products produced using DNA from cells cultured in ~~the~~ conditions excluding ~~any-of~~ any of IL-4, CD40L, TGF  $\beta$  ~~or-and~~ and cycloheximide; as a template. Lane 2 shows the result of Southern hybridization ~~against-of~~ against  
25 PCR products produced using DNA from cells cultured in the presence of cycloheximide only; as a template. Lane 3 shows the result of hybridization ~~against-of~~ against PCR products produced using DNA from cells cultured in the presence of IL-4, CD40L, and TGF  $\beta$ ; as a template. Lane 4 shows the result of hybridization against PCR products produced using DNA from cells cultured in the presence of IL-4, CD40L, TGF  $\beta$  and cycloheximide; as a  
30 template.

FIG. 2 is a pair of photographs showing the production ~~state~~ of DNA including an Sa sequence looped ~~put without~~ by class switch recombination, amplified by PCR using DNA derived from mouse B cell clone CH12F3-2 cultured in various conditions.

FIG. 2(a) shows the ~~electrophoretic-electrophoresis state~~ of ~~DNA from~~ DNA including an Sa sequence looped out ~~with~~ by class switch recombination in mouse B cell clone CH12F3-2 cultured in ~~the~~ various conditions, stained with ethidium bromide.

Lanes 1 and 6 show the ~~electrophoretic-electrophoresis state~~ of marker DNAs. Lane 2 shows the ~~electrophoretic-electrophoresis state~~ of PCR products produced using DNA from cells cultured in ~~the~~ conditions excluding IL-4, CD40L, TGF  $\beta$  ~~or~~ and cycloheximide, as a template. Lane 3 shows the ~~electrophoretic-electrophoresis state~~ of ~~DNA~~ PCR products producing using DNA from cells cultured in the presence of cycloheximide only, as a template. Lane 4 shows the ~~electrophoretic-electrophoresis state for~~ of PCR products produced using DNA from cells cultured in the presence of IL-4, CD40L and TGF  $\beta$ , as a template. Lane 5 shows the ~~electrophoretic-electrophoresis state for~~ of PCR products produced using DNA from from cells cultured in the presence of IL-4, CD40L, TGF  $\beta$ , and cycloheximide, as a template.

FIG. 2(b) shows the result of Southern hybridization ~~for~~ of DNA including an Sa sequence looped out with class switch recombination, amplified by PCR using DNA derived from mouse B cell clone CH12F3-2 cultured under ~~the~~ various conditions.

Lane 1 shows the result of hybridization ~~against~~ of a PCR products produced using DNA from cells cultured under ~~the~~ conditions excluding ~~any one of~~ IL-4, CD40L, TGF  $\beta$  ~~or~~ and cycloheximide, as a template. Lane 2 shows the result of hybridization against a PCR products produced using DNA from cells cultured in the presence of cycloheximide only, as a template. Lane 3 shows the result of hybridization against a PCR products produced using ~~the~~ DNA from cells cultured in the presence of IL-4, CD40L, and TGF  $\beta$ , as a template. Lane 4 shows the result of hybridization against a PCR products produced using DNA from cells cultured in the presence of IL-4, CD40L, TGF  $\beta$  and cycloheximide, as a template.

FIG. 3 shows the result of Northern blotting using a radiolabeled cDNA fragment encoding a ~~radiolabeled~~ 23C9 (AID) protein as a probe, against mRNA derived from mouse B cell clone CH12F3-2 cultured under the various conditions.

Lane 1 shows the result of blotting against mRNA from cells cultured in the conditions excluding ~~any one of~~ IL-4, CD40L, ~~or~~ TGF  $\beta$  ~~or~~ and cycloheximide. Lane 2 shows the result of blotting against mRNA from cells cultured in the presence of cycloheximide, only. Lane 3 shows the result of blotting against mRNA from cells cultured in the presence of IL-4, CD40L, and TGF  $\beta$ . Lane 4 shows the result of blotting against mRNA from cells cultured in the presence of IL-4, CD40L, TGF  $\beta$ , and cycloheximide.

FIG. 4 shows the result of Northern blotting using a radio-labeled cDNA fragment coding 23C9 (AID) protein as a probe against mRNA derived from mouse B cell clone CH12F3-2 cultured in the various conditions.

Lane 1 shows the result of blotting against mRNA from cells cultured in the conditions excluding IL-4, CD40L, TGF  $\beta$  ~~or~~ and cycloheximide. Lane 2 shows the result of blotting against mRNA from cells cultured in the presence of cycloheximide, only. Lane 3 shows the result of ~~the~~ blotting against mRNA ~~form~~ from cells cultured in the presence of IL-4, CD40L, and TGF  $\beta$ . Lane 4 shows the result of ~~the~~ blotting against mRNA ~~form~~ from cells cultured in the presence of IL-4, CD40L, TGF  $\beta$ , and cycloheximide.

FIG. 5 shows the homology between an amino acid sequence of mouse AID protein (SEQ ID NO:2) and that of mouse APOBEC-1 (SEQ ID NO:36).

An amino acid in a closed box shows an identical amino acid. A region in an open box indicates a cytidine deaminase motif. An amino acid with an asterisk (\*) or an arrow indicates an amino acid conserved among APOBEC-1 proteins derived from rat, mouse, rabbit, and human.

FIG. 6 shows a phylogenic tree of various enzymes belonging to a cytosine nucleoside /-nucleotide deaminase family, prepared based on cytidine deaminase motif.

FIG. 7 shows a photograph indicating the electrophoretic state for AID-GST fusion protein in the molecular weight analysis by the gel electrophoresis and silver staining method.

Lane 1 shows the electrophoretic state for a marker molecule. Lane 2 shows the electrophoretic state for various proteins included in extracts from wild type *Escherichia coli* DH 5 $\alpha$ . Lane 3 shows the electrophoretic state for purified AID-GST fusion protein.

FIG. 8 shows the electrophoretic state for AID-GST fusion protein by Western blotting using anti-AID protein peptide antibody.

Lane 1 shows the electrophoretic state for various proteins included in the extract from wild type *E. coli* DH5 $\alpha$ .

Lane 2 shows the electrophoretic state for purified AID-GST protein.

FIG. 9 shows a cytidine deaminase activity depending on the concentrations of AID  
5 proteins.

FIG. 10 shows the inhibitory effect of tetrahydrouridine which is an inhibitor specific to cytidine deaminase on a cytidine deaminase activity in AID protein.

FIG. 11 shows the inhibitory effect of each of 1,10-*o*-phenanthroline which is a zinc-chelating agent, and 1,7-*o*-phenanthroline which is an inactivated isomer thereof on a  
10 cytidine deaminase activity in AID protein.

FIG. 12 is a photograph indicating expression state for mRNA of AID in various tissues in mouse, analyzed by Northern blotting method.

FIG. 13 is a photograph indicating the expression state for mRNA of AID in various lymphatic tissues in mouse, analyzed by RT-PCR method.

FIG. 14 is the photograph showing expression state for mRNA of AID as over time  
15 goes, in activated mouse B cell clone CH12F3-2, analyzed by Northern blotting method.

FIG. 15 a photograph showing expression state for mRNA of AID in mouse B cell clone CH12F3-2 stimulated with cytokine in various combinations, analyzed by Northern blotting.

FIG. 16 shows a photograph indicating expression state for mRNA of AID in mouse  
20 spleen B cells, stimulated with stimulants in various combinations, analyzed by Northern hybridization method.

FIG. 17 is a photograph indicating expression state for mRNA of AID in splenocytes derived from mice immunized with sheep red blood cells, analyzed by Northern blotting  
25 analysis.

FIG. 18 shows expression state for mRNA of AID in splenocytes derived from mice immunized with sheep red blood cells, analyzed by RT-PCR.

FIG. 19 is a photograph indicating localization of expression for AID mRNA in splenocytes derived from a normal mouse or a mouse immunized by sheep red blood cells,  
30 specifically, analyzed by in situ hybridization.

FIGs. 19 (A) and (D) indicate the result in the hybridization using a sense AID probe. FIGs. 19 (B) and (E) show localization for AID mRNA expression in hybridization using an antisense-AID probe. FIGs. 19 (C) and (F) show localization of germinal center in staining test by FITC-labeled PNA. FIGs 19 (A), (B), and (C) indicate the result in the test using  
5 spleen tissues derived from normal mouse (before the immunization of sheep red blood cells). FIGs. 19 (D), (E), and (F) show the results of the examination using spleen tissue slices prepared 5 days after immunizing a mouse with sheep red blood cells.

FIG. 20 is a photograph showing the localization of expression for AID mRNA in spleen tissue and payer's patch tissue, respectively, derived from a normal mouse or from a  
10 mouse immunized with sheep red blood cells, respectively, analyzed by in situ hybridization.

FIGs. 20 (A), (D), and (G) show the results in the hybridization using a sense AID probe. FIGs. 20 (B), (E), and (H) show the localization of the expression for AID mRNA in hybridization using an antisense AID probe. FIGs. 20 (C), (F), and (I) show the localization of germinating center in the staining test by FITC-labeled PNA. FIGs. 20 (A), (B), and (C)  
15 show the result of the examination using spleen tissues derived from a normal mouse (before immunization by sheep red blood cells). FIGs. 20 (D), (E), and (F) show the results of the examination using spleen tissue slices prepared 5 days after immunization of a mouse with sheep red blood cells. FIGs. 20 (G), (H), and (I) show the results of test using payer's patch prepared 5 days after the immunization of a mouse with sheep red blood cells.

20 FIG. 21 schematically shows relative locations of partial nucleotide sequences of human genomic DNA coding human AID protein, which was amplified by PCR using various pairs of primers.

FIG. 22 schematically shows a degree of homology between an amino acid sequence of mouse AID protein (SEQ ID NO:2) and that of human AID protein (SEQ ID NO:8). The  
25 parts with a closed box are cytidine and deoxycytidylate deaminase zinc-binding region which is an AID protein active region.

FIG. 23 schematically shows the structure of human genomic DNA including a gene coding human AID protein. One to five shows exon 1, exon 2, exon 3, exon 4, and exon 5,  
respectively.

30 FIG. 24 is a photograph indicating the expression state for human AID mRNA in various types of human tissues, analyzed by RT-PCR.

FIG. 25 is a photograph indicating a location (localization) of human AID gene on human chromosome, analyzed by Fluorescence in situ hybridization (FISH) method.

Two points at the tips of arrows show 12p13, where the human AID gene exists  
located.-

5 Like reference symbols in the various drawings indicate like elements.

### DETAILED DESCRIPTION OF THE INVENTION

[00195] The invention is illustrated in details by the following Examples, but is not restricted to the embodiments described in the Examples.

10 [00196] Example 1: Culture of ~~for~~ mouse B cell clone CH12F3-2 and confirmation of properties

[00197] Mouse B cell clone CH12F3-2 ~~occurring that undergoes~~ class switch recombination (CSR) from IgM to IgA, several hours after ~~the~~ stimulation by with IL-4, TGF- $\beta$ , and CD40L, previously isolated by the present inventors, was cultured in the same  
15 manner as in the previous report (Immunity, 9:1-10, 1998; Curr. Biol., 8:227-230, 1998; Int. Immunol, 8:193-201, 1996).

[00198] When ~~the~~ CH12F3-2 ~~is~~ cells were stimulated by with IL-4, TGF- $\beta$ , and CD40L, a circular DNA including an S region (switch region) looped out by class switch recombination was detected several hours after the stimulation.

20 [00199] The following manipulation was conducted according to the previously reported method (Curr. Biol., 8:227-230, 1998).

[00200] ~~The CH12F3-2 B cells CH12F3-2, either~~ stimulated by IL-4, TGF- $\beta$ , and CD40L or, and that which was not unstimulated, were cultured for 6 hours in the presence or absence of cycloheximide (200 ng/ml) which is a protein synthesis inhibitor, respectively.  
25 Genomic DNA was extracted from each cell, and PCR was conducted with the DNA as a template ~~by following the standard methods~~ to amplify circular DNA including an S $\mu$  sequence and an S $\alpha$  sequence. PCR was conducted by using a pairs of primers,  $\alpha$ F1 and  $\mu$ R3, or and the other PCR was conducted by using a pair of primers,  $\alpha$ F1 and  $\mu$ R3.

[00201] As a control, genomic DNA encoding glyceraldehyde-3-phosphate  
30 dehydrogenase (GAPDH) was amplified by PCR.

[00202] The PCR products ~~was were~~ subjected to ~~the~~ gel electrophoresis ~~by and~~ ethidium bromide staining. FIG. 1(a) and FIG. 2(a) show the results.

[00203] To confirm the presence or absence of the amplification of a circular DNA including the looped-out S region, Southern hybridization was conducted against the PCR product ~~sby~~ using a mouse S $\alpha$  region gene ~~for as~~ a hybridization probe, according to ~~the~~ standard methods (L. Sambrook E. F., Tom Maniatis., Second edition, Ed. Molecular Cloning (Nolan, C., Ed.) Cold Spring Harbor, 1989). ~~As an S $\alpha$  gene, a~~ 1,155 bp DNA fragment obtained by digesting 10 kb EcoRI cleaved fragment IgH703 with Hind III and EarI was used as an S $\alpha$  gene (Genbank #D11468, DNA No. 1993-3148) (J, Biol. Chem., Vol. 268, p. 4651-4665). FIGs. 1(b) and 2(b) show the results.

[00204] It has been shown that mouse B cell CH2F3-2 produces the looped-out DNA containing the S $\alpha$  sequence with the class switch recombination ~~by the~~ after stimulation with cytokines, and the production of the DNA is inhibited by the presence of cycloheximide. This result suggested that the occurrence of class switch recombination of an immunoglobulin gene ~~needs a novel synthesis of a protein requires de novo protein synthesis~~ in the very early stages ~~after the~~ stimulation, and ~~the that~~ protein synthesis is deeply involved in the induction of ~~the~~ class switch recombination.

[00205] Example 2: Identification of a gene ~~which whose~~ expression is improved in mouse B cell CH12F3-2 stimulated by cytokines.

[00206] A gene which is presumably expressed in the early stage after mouse B cell clone CH12F3-2 is stimulated, and presumably plays a role ~~of in~~ introducing class switch recombination of an immunoglobulin gene, was attempted to be isolated ~~from the~~ CH12F3-2 cells by the suppression subtract hybridization (SSH) (Proc. Natl. Acad. Sci. USA, 93:6025-6030, 1996; Anal. Biochem., 240:90-97, 1996) using the inhibitory PCR effect (Nucleic Acids Res., 23:1087-1088, 1995).

[00207] A cDNA library necessary for subtraction cloning was prepared ~~by using a~~ PCR-Select Subtraction Kit (CLONTECH, Catalogue NO: K1804-1) by following the instruction manual ~~supplemented supplied~~ with the kit ~~in the experimental manipulation~~.

[00208] PolyA<sup>+</sup>-RNA was isolated from each of: mouse B cell clone CH12F3-2 stimulated with IL-4, TGF- $\beta$  and CD40L for 5 hours, the same cells stimulated with the

cytokines for 12 hours, and the cells which were not stimulated, by following the reported method (Nucleic Acids Res., 26:911-918, 1998) and treated with DNaseI to eliminate any contaminating genomic DNA ~~which may be mixed~~. Then cDNA was prepared based on each polyA<sup>+</sup>-RNA sample using reverse transcriptase according to the standard method. Each  
5 cDNA prepared from mouse B cell clone CH12F3-2, treated with the above cytokines for 5 or 12 hours, was mixed ~~with same~~ in equimolar mole amounts to be used as a tester cDNA. On the other hand, cDNA derived from unstimulated cells was used as a driver cDNA.

[00209] Subtraction was conducted by adding the driver cDNA into the tester cDNA according to the ~~above~~ above-referenced previous report and the ~~experimental manipulation~~  
10 kit instruction manual. The efficiency of subtraction was monitored by adding a small amount (1:1000 mole ratio) of  $\Phi$ -X174 phage DNA cleaved at the restriction enzyme site Hae III, as a control, into the tester cDNA. After the subtraction, the phage DNA was concentrated to a mole ratio of about 100 times.

[00210] The subtracted cDNA was cloned into the T-vector (Promega) according to the  
15 standard method to prepare a plasmid library. In the same manner as in the previous report, 2000 colonies in the library were screened by the differential hybridization method (Nucleic Acids Res., 26:911-918, 1998; Medical immunity, 29:17, p. 451-459, 1997). Each of the above tester cDNA and driver cDNA was radiolabeled to be used for hybridization. Clones including  $\Phi$  X174 phage DNA were selected by hybridizing  $\Phi$  X174 phage DNA with a  
20 replicant filter.

[00211] One hundred fifteen clones emitting a stronger signal than the radio-labeled driver cDNA probe against ~~radio-labeled~~ radiolabeled tester cDNA probe were identified and a nucleotide sequence of each clone was determined by using a DNA sequencer.

[00212] Northern blotting was conducted against mRNA obtained from mouse B cell  
25 clone CH12F3-2 stimulated with IL-4, TGF- $\beta$  and CD40L or the same cell lines unstimulated, using the radio-labeled DNA inserted into the each clone as a probe, according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition, Ed. Molecular Cloning (Nolan, C., Ed.), Cold Spring Harbour, 1989). As a result, the enhanced expression corresponding to the stimulation with the above cytokines was observed in 23 ~~among out of~~  
30 115 clones. Gene fragments coding 7 different types of proteins, including genes coding ~~the~~ 3 kinds of known proteins and 4 kinds of novel proteins were found to be inserted into the 23



clones. Specifically, the expression of the 7 kinds of genes ~~were~~was found to be enhanced in mouse B cell clone CH12F3-2 by the stimulation with IL-4, TGF- $\beta$  and CD40L.

**[00213]** <The known proteins>

- 5 ABCD-1/MDC (8 clones)  
IFN $\gamma$  receptor (2 clones)  
I-a (MHC class II) (1 clone)

**[00214]** <Novel proteins>

- 10 23C9 (3 clones)  
15B11 (7 clones)  
8B9 (1 clone)  
16A9 (1 clone)

- 15 **[00215]** As it has been known that the expression of the above I-a gene and ABCD/MDC gene is enhanced by stimulating mouse spleen B cell with IL-4 and CD40L, it was confirmed that the subtraction cloning was effectively conducted (J. Exp. Med., 188:451-463, 1998; Immunity, 5:319-330, 1996).

- 20 **[00216]** Example 3: Expression of mRNA for a novel protein 23C9 in mouse B cell clone CH12F3-2

- [00217]** The degree of enhanced expression of the gene coding for a novel protein 23C9 in mouse B cell clone CH12F3-2 stimulated with IL-4, TGF- $\beta$  and CD40L was analyzed according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition, Ed. Molecular Cloning (Nolan, C., Ed.), Cold Spring Harbour, 1989) by Northern blotting.

**[00218]** Mouse B cell clone CH12F3-2 was cultured in the presence of one of the following reagents for 12 hours.

- (1) IL-4, TGF- $\beta$  and CD40L only.  
30 (2) Cycloheximide which is a protein synthesis inhibitor (200 ng/ml), only  
(3) IL-4, TGF- $\beta$  and CD40L as well as Cycloheximide (200 ng/ml)

[00219] Northern blotting was conducted against mRNA (10 µg for each group) obtained in the same manner as ~~the previously~~ reported (Nucleic Acid Res., 26:911-918, 1998) from each group of treated cells using a radio-labeled cDNA fragment (1,020 bp) coding- for a novel protein 23C9, obtained in the above Example, according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition, Ed. Molecular Cloning (Nolan, C., Ed.), Cold Spring Harbour, 1989).

[00220] As a control ~~examination~~experiment, Northern blotting was conducted for mRNA derived from B cell clone CH12F3-2 cultured without any one of the above cytokines, or cycloheximide.

[00221] The amount of mRNA to be electrophoresed was adjusted using the amount of mRNA ~~in~~ for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) as an index. DNA amplified by RT-PCR using a GF primer and a GR primer was used as a probe for ~~blotting~~blots of GAPDH mRNA (Location of nucleotides: 566-1016, Genbank U5299) (Immunity, 9:1-10, 1998).

[00222] FIGs. 3 and 4 show the results.

[00223] The expression of mRNA for a novel protein 23C9 was extremely strong in mouse B cell clone CH12F3-2 stimulated with IL-4, TGF-β and CD40L, while the expression in unstimulated cells was extremely weak. Expression of the mRNA in the stimulated cells was inhibited by the presence of a protein synthesis inhibitor. Moreover, in the stimulated cells, two bands indicating the expression of mRNAs comprising different nucleotide lengths were detected.

[00224] Expression of mRNA for a novel protein 23C9 in each of the following mouse cell lines, which do not originally ~~comprise~~ undergo class switch recombination, was examined by Northern blotting in the same manner as in the above.

[00225] B cell lines (lyD9, BA/F3, 70Z/3, WEHI231); T cell lines (EL-4, 2B4); myeloma cell lines (X63, HEHI-3); ~~Fibroblast~~ fibroblast lines (L929, NIH3T3); ~~the and~~ other cell lines (F2, P815, ST2).

[00226] The expression of mRNA for the novel protein 23C9 was not observed in any of these cells.

[00227] Example 4: Cloning of a full length cDNA encoding a novel protein 23C9.

[00228] Four different positive clones were obtained by screening a cDNA library (Nucleic Acids Res., 26:911-918, 1998) prepared from mouse B cell clone CH12F3-2 stimulated with IL-4, TGF- $\beta$ , and CD40L, using a cDNA fragment (1,020 bp) encoding the novel protein 23C9, obtained in the above Example, as a probe. ~~A~~The nucleotide sequence of each clone was determined by using a DNA sequencer according to the standard method.

[00229] One clone comprises a 1.2 kb nucleotide length and a single reading frame (~~OPFORF~~) with 1 polyadenylation site. The other 3 clones comprise a 2.4 kb nucleotide length and 2 polyadenylation sites. ~~A~~The nucleotide sequence ~~in of the~~ first 1.2 kb ~~part~~ at the 5' side ~~in end of~~ the latter clones was identical to that of the nucleotides ~~in of the~~ 1.2 kb DNA in the former (SEQ ID NO:1).

[00230] Two different mRNA transcripts detected in Northern blotting in the above Example (FIGs. 3 and 4) were predicted to correspond to transcripts for each of the above 1.2 kb and 2.4 kb, transcribed ~~using until~~ the polyA site at the 3' end and the polyA site at the 5' end.

[00231] A cDNA fragment encoding the novel protein 23C9 used as a probe in the above (1,020 bp) was found to ~~be have~~ a nucleotide sequence of from 847 to 1866 in the full length cDNA of 23C9.

[00232] A nucleotide sequence near a first initiation codon in each cDNA was fit to Kozak's rule (Nucleic Acids Res., 15:8125-8148, 1987). In the 2.4kb cDNA, ATTTA, which is a motif capable of mediating quick degradation of mRNA (Blood, 83:3182-3187, 1994), was present in the 2 sites in the 3' untranslated ~~ional~~ region ~~in the 3' side~~.

[00233] An open reading frame (ORF) of cDNA encoding ~~a the~~ novel protein 23C9 consisted of 198 amino acids with the expected molecular weight of about 24 kDa (SEQ ID NO:2). ~~As a result of homology searching with known proteins by in a database, it was discovered that the~~ amino acid sequence ~~of of the~~ ORF of the novel protein 23C9 comprised 34% amino acid homology with the apolipoprotein B mRNA editing enzyme, catalytic polypeptide-1 (APOBEC-1) (Science, 260:1816-1819, 1993; J. Biol. Chem., 268:20709-20712, 1993). GenBank and EMBL were used as the DNA data bases. SwissPlot was used as the protein database. The BLAST program (J. Mol. Biol., 215:403-410, 1990) and the FASTA program (Proc. Natl. Acad. Sci. USA., 85:2444-2448 1988) were used ~~for to~~

search the databases-search.

FIG. 5 shows an amino acid sequence of the ORF of the novel protein 23C9 and an alignment between the sequence and that of mouse APOBEC-1 amino acid sequence.

[00234] ~~As a result of motif~~ Motif searching ~~on~~-online using PROSITE (Nucleic Acids Res., 11:2013-2018, 1992), indicates that the APOBEC-1-like novel protein 23C9 comprises a cytidine/deoxycytidine deaminase motif which is conserved in ~~a~~the amino acid sequences of ~~a~~ proteins belonging to ~~to~~ the cytosine nucleoside/-nucleotide deaminase family, which ~~constructs~~ is a large family and is an activation site of a deaminase activity. ~~A~~The cytosine nucleoside/nucleotide deaminase family is classified into RNA editing deaminases, cytidine/deoxycytidylate deaminases, and CMP/dCMP deaminases based on ~~the~~ substrate specificity and homology in the activation sites (Cell, 81:187-195, 1995).

[00235] ~~A phylogenie~~ phylogenetic tree was prepared based on the alignment ~~among~~ of regions in APOBEC-1, which is an RNA editing deaminases, cytosine nucleoside deaminase, cytosine nucleotide deaminases, and the cytidine deaminase motif in the novel protein 23C9. ~~The sequences in~~ of the known proteins used for the comparison were obtained from GenBank, as follows.

Human derived nucleoside deaminase: L27943

Mouse derived nucleoside deaminase: AA388666

*S. subtilis* derived nucleoside deaminase: U18532

*E. coli* derived cytidine deaminase: X63144

Rabbit derived APOBEC-1: U10695

Human derived APOBEC-1: L25877

Rat derived APOBEC-1: U10695

Mouse derived APOBEC-1: U21951

T2/T4 phage derived nucleotide deaminase: J05172

Human derived nucleotide deaminase: L12136

*S. cerevisiae* derived nucleotide deaminase: U10397

[00236] FIG. 6 shows the result. ~~Cytidine~~ The cytidine deaminase motif in the novel protein 23C9 was ~~rather~~ relative more closely related to a subgroup of RNA editing deaminases than to subgroups of nucleoside deaminase and nucleotide deaminase.

[00237] On the other hand, a leucine-rich region existing at the C-terminus of APOBEC-1 is thought to be important for protein-protein interaction (Proc. Natl. Acad. Sci. USA., 91:8522-8526, 1994; J. Biol. Chem., 269:21725-21734, 1994). The novel protein 23C9 also comprised a leucine-rich region at the C-terminus. Four leucines in the region of  
5 23C9 were conserved in the leucine rich regions of APOBEC-1 in rabbit, rat, mouse and human.

[00238] It has been known that Phe66, Phe87, His61, Glu63 and Cys93 are essential for binding of APOBEC-1 to RNA, and all these amino acid residues were conserved in the primary structure of 23C9 (Trends Genet., 12:418-424, 1996; Cell, 81:187-195, 1995; J. Biol.  
10 Chem., 270:14768-14775 1995; J. Biol. Chem., 270:14762-14767, 1995). From this fact, 23C9 protein is predicted to comprise an RNA editing deaminase activity.

[00239] Moreover, cytidine deaminases derived from APOBEC-1 and *E. coli* (ECCDA) are known to comprise a pseudoactive site domain at the C-terminus, and the 23C9 protein also ~~comprised~~ comprises a pseudoactive site domain, the same as is in the  
15 APOBEC-1. This indicates that 23C9 protein is more closely related ~~relative~~ to APOBEC-1 and ECCDA than to deaminase proteins in the other groups.

[00240] From these facts, the novel protein 23C9 was named activation-induced cytidine deaminase (AID). The novel protein 23C9 was called AID hereafter.

20 [00241] Example 5: Preparation of the AID-GST fusion protein

[00242] The cDNA coding ~~a~~ the full length AID cloned in the above Example was amplified by PCR with a pair of primers, AID-138 (SEQ ID NO:3) and AID-161 (SEQ ID NO:4), a pair of primers, AID-118 (SEQ ID NO:5) and AID-119 (SEQ ID NO:6), using ~~and~~ Taq Polymerase ~~by~~ following the standard method. As there is an intron between AID-118  
25 and AID-119, a PCR product derived from AID genomic DNA can be easily distinguished.

[00243] The obtained PCR product was subcloned ~~to~~ into the pGEX4T1 vector (Pharmacia) according to the standard method. A nucleotide sequence of the vector was determined and the absence of point mutations derived from the use of Taq polymerase in the full length AID cDNA cloned ~~to~~ into the vector was confirmed.

30 [00244] *E. coli* DH5 $\alpha$  was transformed with the vector according to the standard method. The obtained transformants were cultured, and a full length AID cDNA was

expressed as a fusion protein with glutathione S-transferase (GST). The AID-GST fusion protein was extracted in the same manner as in the previous report, and purified using glutathione agarose affinity chromatography (J. Biol. Chem., 270:14768-14775 1995).

[00245] A-The molecular weight of the purified AID-GST fusion protein was analyzed

5 by following the standard method using 10% SDS-PAGE and silver staining. A protein extracted from wild type *E. coli* DH5a was used as a control. FIG. 7 shows the results.

[00246] As expected, the fusion protein was detected as a band comprising a molecular weight of about 49 kDa. Minor bands detected under the about 49 kDa were thought to be decomposed proteins, frequently generated in the purification process in  
10 general.

[00247] A molecular weight of the purified AID-GST fusion protein was analyzed by the Western blotting method according to the standard method (Genomics, 54:89-98, 1998).

Anti-AID protein antibody to be used for the assay was prepared by immunizing a commercial rabbit for the experiment with multiple antigen peptides including synthetic  
15 peptides corresponding to amino acids ~~NONos~~: 116 to 132 of the AID protein of the present invention (Proc. Natl. Acad. Sci. USA., 85:5409, 1988).

[00248] FIG. 8 shows the results.

[00249] Example 6: Cytidine deaminase activity of the AID protein

20 [00250] A cytidine deaminase activity of AID was measured by the same method as in the previous report (J Biol. Chem. 270:14768-14775, 1995).

[00251] The purified AID-GST fusion protein prepared in the above (2, 4, 6, 8, 10, 20, 40, 60, 100, 200, 300, 400, and 600 ng) was incubated in the buffer (pH 7.5, the total amount 10 µl) containing 45mM Tris with 3.3 µCi [<sup>3</sup>H] deoxycytidine (24.8 Ci/mmol, DuPont) and  
25 250 µM cytidine for 2 to 4 hours. The reaction was terminated by adding deoxycytidine (2 µl of 10 µg/ml) and deoxyuridine (2 µl of 10 µg/ml). Insoluble substances were removed by centrifugation, and the reaction mixture (4 µl) was subjected to the polyethylene imine-cellulose thin layer chromatography plate (VWR). The plate was developed in isopropyl alcohol/-10% HCl (7:2 v/v). The plate was exposed to ultraviolet light (254 nm) for  
30 visualization and bands corresponding to deoxycytidine and deoxyuridine were collected,

and added to Ultima Gold scintillation solution to be quantified by liquid scintillation photometer (Packard)

[00252] FIG. 9 shows the results. ~~As a result~~The AID protein showed ~~the a~~  
concentration-dependent cytidine deaminase activity ~~depending on the concentrations~~.

5 [00253] ~~An~~The inhibitory effect of tetrahydrouridine (THU; 0 to 40  $\mu$ M)  
(Calbiochem, USA), which is an inhibitor specific to cytidine deaminase, on the cytidine  
deaminase activity in the AID-GST fusion protein (300 ng) was measured by the same  
method described above.

[00254] FIG. 10 shows the results. The cytidine deaminase activity of AID protein was  
10 inhibited ~~dependently~~depending on the concentrations of THU.

[00255] ~~Each~~The inhibitory effect of 1,10-*o*-phenanthroline (0 to 20 mM), which is a  
zinc-chelating agent, and its inactive ~~form from~~ isomer 1,7-*o*-phenanthroline (0 to 20 mM),  
on the cytidine deaminase activity ~~in of~~ the AID-GST fusion protein was measured in the  
same manner as described below.

15 [00256] FIG. 11 shows the results. The cytidine deaminase activity of AID protein was  
inhibited by 20 mM 1,10-*o*-phenanthroline by about 91%. 1,7-*o*-phenanthroline, which is the  
inactive isomer, only inhibited about 13%. These results indicate that the AID protein is a  
zinc-dependent cytidine deaminase, ~~same as~~similar to APOBEC-1.

20 [00257] Example 7: Avidity of AID protein with AU-rich RNA

[00258] A recombinant APOBEC-1 binds to ~~Au~~AU-rich RNA (Trends Genet., 12:418-  
424, 1996; Cell, 81:187-195, 1995; J. Biol. Chem., 270:14768-14775, 1995; J. Biol. Chem.,  
270:14762-14767, 1995), and progresses RNA editing for apoB in the presence of chicken  
extract including co-factor.

25 [00259] Since the AID protein has a functional cytidine deaminase activity as well as a  
structural similarity ~~with to~~ APOBEC-1, to examine ~~an~~ RNA editing activity in the AID  
protein, avidity to AU-rich RNA (5-AU) and apoB RNA (which are RNA substrates for  
APOBEC-1) was examined.

[00260] The AID protein did not show avidity to AU-rich RNA (5-AU) in the gel  
30 retardation assay. In an *in vitro* apoB RNA assay, conversion from cytidine (C) to uridine (U)  
was not observed.

[00261] Example 8: Expression distribution of AID mRNA in tissues

[00262] The expression of AID mRNA in each tissue was examined by Northern blotting according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition, Ed. Molecular Cloning (Nolan C., Ed.), Cold Spring Harbour, 1989; Experimental Medicine, Suppl., "Genetic Engineering Hand Book", published by Yodosha, p. 133-140, 1992).

[00263] PolyA<sup>+</sup>-RNA (2 µg each) obtained from cells derived from each tissue in mice (muscle, spleen, lung, heart, lymph node, brain, kidney, thymus, testis, liver) according to the previous report (Nucleic Acids Res., 26:911-918, 1998) was used as a sample. Radiolabeled cDNA fragments (1,020 bp) encoding AID (23C9) obtained in the previous Examples was used as a probe for blotting polyA<sup>+</sup>-RNA.

[00264] As a control, mRNA of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was blotted in the same manner. ~~As~~ As a probe for blotting GAPDH mRNA, DNA amplified by PCR using GP primer and GR primer was used. (Nucleotide location: 566-1016, Genbank, U52599) (Immunity, 9:1-10, 1988).

[00265] FIG. 12 shows the results.

[00266] ~~As a~~ The results show that, AID mRNA was strongly expressed in mesenteric lymph node. In addition, weak expression was observed in spleen.

[00267] Example 9: Expression of AID mRNA in various lymphatic tissues.

[00268] The expression of AID mRNA in each lymphatic tissue was analyzed by RT-PCR according to the standard method (Immunity, 9:1-10, 1998).

[00269] cDNA was prepared according to the standard method using polyA<sup>+</sup>-RNA obtained from cells derived from various lymphatic tissues (Payer's patch, mesenteric lymph node, axillary lymph node, spleen, bone marrow, thymus) in the same manner as in the previous report (Nucleic Acids Res., 26:911-918, 1998), for mRNA as a sample, as a template. AID cDNA and GAPDH cDNA were amplified using the obtained cDNA as a template. ~~The~~ A first pair of primers, AID-138 (SEQ ID NO:3) and AID-161 (SEQ ID NO. 4) ~~in the~~ as above, a second pair of primers AID-118 (SEQ ID NO:5) and AID-119 (SEQ ID NO:6), and Taq polymerase were used for PCR of AID cDNA. As there is an intron between



AID-118 and AID-119, a PCR product derived from the AID genomic DNA sequence can be easily distinguished.

[00270] FIG. 13 shows the results.

[00271] AID cDNA was detected in all lymphatic tissues except for thymus. In particular, ~~the~~ obvious expression was observed in peripheral lymphatic organs, such as lymph node or Payer's patch. On the other hand, ~~the~~ expression in primary lymphatic organs was weak in comparison with that in the peripheral lymphatic organs.

[00272] Example 10: Expression of AID mRNA ~~as over~~ time ~~goes in~~ activated mouse B cell clone CH12F3-2

[00273] Expression of AID mRNA ~~as over~~ time ~~goes in~~ activated mouse B cell clone CH12F3-2 stimulated with IL-4, TGF- $\beta$ , and CD40L for 0 to 60 hours was analyzed by Northern blotting according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition Ed. Molecular Cloning (Nolan, C., Ed.). Cold Spring Harbor, 1989).

[00274] Mouse B cell clone CH12F3-2 was cultured in the presence of IL-4, TGF- $\beta$ , and CD40L for ~~the~~ various periods (0, 3, 5, 12, 24, 36, 48, or 60 hours).

[00275] Northern blotting was conducted against mRNA (10  $\mu$ g in each group) obtained from each culture group in the same manner as in the previous report (Nucleic Acids Res., 26:911-918, 1998) using a radiolabeled cDNA fragment encoding AID (23C9) obtained in the previous Examples, as a probe, according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition Ed. Molecular Cloning (Nolan, C., Ed.). Cold Spring Harbor, 1989).

[00276] The amount of mRNA to be gel-electrophoresed was adjusted by using mRNA of GAPDH as an index. DNA amplified by RT-PCR using a GF primer and a GR primer was used as a probe for blotting GAPDH mRNA (Nucleotide location: 566-1016, Genbank U52599) (Immunity, 9:1-10, 1998).

[00277] FIG. 14 shows the results.

[00278] It was shown that ~~T~~the expression of AID mRNA in mouse B cell clone CH12F3-2 was too ~~small~~ low to be detected without ~~the~~ stimulation by cytokines, but ~~that the~~ expression was initiated 3 hours after ~~the~~ stimulation by cytokines (described in the above),

was maximum 12 hours after the stimulation (more than about 15 times), and was gradually decreased from 48 hours after the stimulation.

[00279] Example 11: Cytokine specificity ~~to inducing~~ in the induction of expression of AID mRNA in mouse B cell clone CH12F3-2.

[00280] Cytokine specificity ~~to in~~ inducing expression of AID mRNA in mouse B cell clone CH12F3-2 was analyzed by Northern blotting according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition Ed. Molecular Cloning (Nolan, C., Ed.), Cold Spring Harbour, 1989).

[00281] Mouse B cell clone CH12F3-2 was cultured in the presence of various combinations of cytokines (one or more selected from IL-4, TGF- $\beta$ , and CD40-L) for 12 hours. Northern blotting was conducted against mRNA (10  $\mu$ g in each group) obtained from each culture group in the same manner as in the previous report (Nucleic Acids Res., 26:911-918, 1998) using a radio-labeled cDNA fragment (1,020 bp) coding AID (23C9) obtained in the previous Example, according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition Ed. Molecular Cloning (Nolan, C., Ed.). Cold Spring Harbor, 1989).

[00282] The amount of mRNA to be gel-electrophoresed was adjusted by using mRNA of GAPDH as an index. DNA amplified by RT-PCR using a GF primer and a GR primer was used as a probe for blotting GAPDH mRNA (Nucleotide location: 566-1016, Genbank U52599) (Immunity, 9:1-10, 1998).

[00283] FIG. 15 shows the results.

[00284] ~~Expression i~~Induction of AID-mRNA expression ~~was small~~ by solely any one kind of cytokines was small. On the other hand, when 3 kinds of cytokines as described above were used at the same time, ~~the maximum expression~~ maximal induction of AID-mRNA expression was observed.

[00285] As described in the above Example 3, because ~~expression~~ induction of AID mRNA expression was inhibited by cycloheximide, which is an inhibitor ~~for~~ of protein synthesis, it is hypothesized that enhanced expression of AID mRNA ~~needs~~ requires *de novo* protein synthesis.

[00286] Example 12: ~~Expression~~ Induction of AID mRNA ~~Expression~~ in spleen B cells by stimulation

[00287] The presence or absence of expression ~~induction~~ of AID mRNA ~~by~~ after stimulation which may activate B cells and induce class switch recombination of

5 immunoglobulin was examined.

[00288] Spleen B cells ~~was~~ were obtained ~~purified and obtained~~ from BALB/c mouse (6 to 12-week old, Shimizu Experimental Materials (SLC)) and purified according to the standard method. Dead cells and cell fragments were removed by Ficoll density gradient centrifugation after the process of removing T cells. The purified spleen B cells ~~were~~ was cultured for 4 days in the presence of a stimulus in various combinations (one or more selected from IL-4, TGF- $\beta$ , CD40L, and LPS (lipopolysaccharide) in the same manner as in the previous report (Nucleic Acids Res., 26:911-918, 1998). LPS derived from Salmonella typhosa (50  $\mu$ g/ml, Sigma) was used.

[00289] Northern blotting was conducted against mRNA (15  $\mu$ g in each group) obtained from each culture group in the same manner as in the previous report (Nucleic Acids Res., 26:911-918, 1998) using a radio-labeled cDNA fragment encoding AID (23C9) obtained in the previous Example, according to the standard method (L. Sambrook, E. F., Tom Maniatis., Second edition Ed. Molecular Cloning (Nolan, C., Ed.): Cold Spring Harbor, 1989).

20 [00290] The amount of mRNA to be gel-electrophoresed was adjusted by using mRNA of GAPDH and 28S ribosomal RNA as an index. DNA amplified by RT-PCR using a GF primer and a GR primer was used as a probe for blotting GAPDH mRNA (Nucleotide location: 566-1016, Genbank U52599) (Immunity, 9:1-10, 1998).

[00291] FIG. 16 shows the results.

25 [00292] The enhanced expression of AID mRNA by ~~the~~ stimulation with LPS only, ~~or~~ LPS+IL-4, or LPS+TGF- $\beta$  was observed in normal mouse spleen B cells.

[00293] Example 13: Induced expression of AID mRNA *in vivo*

[00294] It was examined whether the ~~expression~~ induction of AID mRNA expression by various stimulations in vitro would also occur *in vivo*.

30

[00295] BALB/c ~~mouse-mice~~ (6 to 12-week old, five individuals in each group, SLC) ~~was-were~~ immunized by intraperitoneally administering sheep red blood cell (SRBC) ( $1 \times 10^8$  cells, Cosmo Bio.). In the living body immunized by SRBC, it has been known that clonal expansion and germinal center formation occur after the immunoresponse, and class switch recombination of an immunoglobulin gene and affinity maturation are caused.

[00296] PolyA<sup>+</sup>-RNA was prepared from splenocytes isolated from spleen excised from each individual before (day 0) and after (day 2, 5 and 13) the immunization.

[00297] The ~~PolyA~~PolyA<sup>+</sup>-RNA (2 µg each) was subjected to ~~the~~Northern blotting using the radiolabeled cDNA fragment (1,020 bp) encoding AID (23C9) as a probe in the same manner as the above Examples. The amount of mRNA to be gel-electrophoresed was adjusted using mRNA of GAPDH as an index in the same manner as in the above Examples.

[00298] FIG. 17 shows the results.

[00299] The minimum amount of expressed AID mRNA was detected before immunization of SRBC (day 0), however, a significant enhancement of expression (about 4 to 5 times) was observed day 5 and day 13 after the immunization.

[00300] Moreover, to analyze in which cell type enhanced expression of AID mRNA occurs, RT-PCR was conducted by the standard method (Immunity, 9:1-10, 1998).

[00301] Red blood cells were removed from splenocytes obtained from spleen which was obtained 5 days after the immunization of SRBC in the same manner as the above, and T cells and non-T cells were separated using nylon fiber (Wako Pure Chemicals) in the same manner as in the previous report (Eur. J. Immunol., 3:645-649, 1973). T cell fractions contained more than 90% of CD3 positive cells, and less than 20% B 220 positive cells.

[00302] ~~T-cell fraction (after removal of B cells) and B-cell fractions were concentrated~~Concentration of T cell fractions (removal of B cells) and concentration of B cell fractions were carried out by MACS method with magnetic beads conjugated to anti-CD19 antibody (Miltenyi Biotech.). B220 positive B cells included in the fraction in which T cells were removed were 5% or less. On the other hand, B220 positive B cells included in the fraction in which CD19 positive cells were concentrated were 60% or more.

[00303] cDNA was prepared by reverse transcriptase according to the standard method using polyA<sup>+</sup>-RNA prepared from each fractionated cell group. AID cDNA and GAPDH cDNA were amplified by PCR using the obtained cDNA as a template. For PCR of AID

cDNA, the previously described pair of primers, AID-138 (SEQ ID NO:3) and AID-161 (SEQ ID NO:4), and the previously described pair of primers, AID-118 (SEQ ID NO:5) and AID-119 (SEQ ID NO:6), as well as Taq polymerase, were used.

[00304] FIG. 18 shows the results.

5 [00305] ~~As a result, i~~In the CD19 positive B cell fraction and non-T cell fraction, amplification of AID cDNA was observed. Specifically, it was demonstrated that enhanced expression of AID mRNA induced by immunization by SRBC occurs in spleen CD19 positive B cells.

10 [00306] Example 14: Localization of AID mRNA expression in lymphatic organs

[00307] It was found that timing of enhanced expression of AID mRNA in spleen is almost consistent-coincident with the initiation of germinal center (GC) formation after immunization of SRBC, from the result of the previous Examples. In this ~~examination experiment, an the~~ precise localization of AID mRNA expression in lymphatic  
15 organs was analyzed using in situ hybridization.

[00308] AID cDNA, cleaved out by digesting the pGEX4T1 vector in which cDNA encoding the AID protein has been subcloned; with EcoRI and XhoI, was subcloned into plasmid pB1uescriptSK (+) (Stratagene). The plasmid was digested with EcoRI or XhoI to obtain linearized plasmid DNA and transcribed into RNA using the plasmid as a template in  
20 the presence of digoxigenin-labeled rUTP (Boehringer-Mannheim) using T3 RNA polymerase or T7 RNA polymerase to prepare ~~each of~~ digoxigenin-labeled antisense probes and sense probes.

[00309] ~~On the other hand, f~~Frozen tissue slices were prepared by immobilizing with paraformaldehyde from each of the spleen and ~~payer's~~ Payer's patch in a normal mouse as a  
25 lymphatic organ sample. A normal mouse was immunized with SRBC in the same manner as in the above Examples, and frozen tissue slices immobilized with paraformaldehyde ~~from spleen~~ were obtained from spleen 5 days after the immunization.

[00310] Hybridization was conducted by applying the digoxigenin-labeled antisense AID probes or sense AID probes to each of the slides furnished with each of immobilized  
30 slices. Hybridized digoxigenin-labeled AID probe was detected using anti-digoxigenin antibody conjugated with alkaline phosphatase. The localization of anti-digoxigenin

antibody conjugated to digoxigenin on the probe was identified by detecting a phosphatase reactant (dark purple color). This analysis was conducted using a light transmission microscope.

[00311] *In situ* hybridization and detection of riboprobes in this examination  
5 experiment were conducted in the same manner as in the previous report (J. Comp. Neurol., 333:398-416, 1993).

[00312] The location of the germinal center in each tissue slice was identified by staining with PNA (Vector) conjugate with FITC and observing with a inflorescent microscope.

10 [00313] FIGs.19 and 20 show the results.

[00314] In the ~~examination~~ experiments using the antisense AID probe, multiple obvious focal signals were observed in spleen tissue slices derived from SRBC immunized mouse (day 5 after the immunization) (FIGs. 19(E) and 20(E)), however, ~~any~~ no signals were ~~not~~ detected in spleen tissue slices derived from mice which were not immunized with SRBC  
15 (FIGs. 19(B) and 20(B)). This result is consistent with the result of Northern blotting obtained in the above Example (FIG. 17). ~~The~~ Existence of a germinal center was observed both in spleen tissue slices derived from SRBC-immunized mouse (day 5 after the immunization) (FIGs. 19(F) and 20(F)) as well as in the normal ~~payer's~~ Payer's patch (FIG. 20(I)), by staining with FITC-labeled PNA. The expression of AID mRNA was found to  
20 localize in the germinal center in ~~the~~ both tissue slices.

[00315] In the ~~examination~~ experiments using the sense AID probe, ~~any signals as background were not detected~~ no background signals were detected in tissue slices of spleen or of ~~p~~ Payer's patch regardless of the presence or absence of the immunization by SRBC.

[00316] This result indicates that the ~~expression~~ induction of AID mRNA expression  
25 occurs ~~in~~ specifically to in germinal center B cells, activated by stimulation with an antigen.

[00317] Example 15: Isolation of human genomic DNA encoding the AID protein  
~~derived from human~~

15-1 Preparation of probes for hybridization

30 [00318] PCR was conducted using an expression vector, prepared by inserting cDNA encoding a full length mouse AID protein, (prepared in Example 5) into a plasmid vector

pGEX4T1, as a template, with a pair of primers (Primer 170: SEQ ID NO:16 and primer 181: SEQ ID NO:179), according to the standard method described in the above.

5 [00319] The ~~obtained~~ resulting PCR product was purified by the standard methods described above and a nucleotide sequence of the purified DNA was determined by ~~the a~~ direct ~~sequence~~ sequencing method to confirm that the purified DNA is the nucleotide sequence encoding a full length of mouse AID protein. This purified DNA was used as a probe for hybridization in the following experiments.

#### 15-2 Screening of human genomic DNA library

10 [00320] The probe prepared in the above was labeled in the same manner as ~~for~~ the radioactive probe in the above Northern hybridization to make a probe radio-labeled by a radioactive isomer.

[00321] Using the labeled probe, a commercial human genomic DNA library (catalogue No. HL1067j; Lot No. 45003; CLONTECH) was screened by ~~the~~ cross  
15 hybridization according to the standard method.

[00322] Washing after ~~the~~ hybridization was conducted twice in 2 X SSC (including 0.1% SDS, ~~under the~~ at room temperature, 10 min), and twice in 2 X SSC (including 0.1% SDS, 65°C, 30 min). Phage DNA was subcloned by purifying phage DNA and inserting  
20 about 22 kb genomic DNA, obtained by cleaving at NotI restriction enzyme site in the phage DNA, into the Not I restriction enzyme site in plasmid pZero-2.1. This plasmid was named 3CpZero.

[00323] A DNA fragment obtained by digesting 3CPZero with PstI was ligated to the PstI site of plasmid pBlueScript KS (Toyobo) and *E. coli* was transformed with this ligated DNA.

25 [00324] Transformants were screened by the colony hybridization using the labeled probe prepared in the above according to the standard method, and multiple positive clones were obtained.

[00325] ~~A~~ The nucleotide sequence of human genomic DNA inserted into each positive clone was analyzed and multiple clones containing genomic DNA of DNA encoding  
30 a human AID protein were identified.

[00326] Among the multiple clones, nucleotide sequences of genomic DNA containing DNA encoding a human AID protein contained in two clones ~~were~~are described in SEQ ID NOs:9 and 10, respectively.

5 [00327] ~~Moreover~~In addition, ~~a~~the nucleotide sequence of genomic DNA including the DNA encoding a human AID protein included in the ~~positive~~other positive clone ~~was~~is shown in SEQ ID NO:35.

[00328] Example 16: Isolation of cDNA encoding a full-length human AID protein and preparation of human AID protein

10 [00329] By comparing ~~a~~the nucleotide sequence of genomic DNA, including ~~a~~the coding region of the ~~obtained~~ human AID protein, with cDNA nucleotide sequence encoding a~~the~~ full-length mouse AID protein (determined in the examples above), a human AID protein encoding region in the human genomic DNA was deduced.

15 [00330] A pair of primers for RACE-PCR was designed based on the deduced nucleotide sequence of the coding region in the human AID protein (Primer 22: SEQ ID NO:18, and primer 25: SEQ ID NO:19).

[00331] RACE-PCR was conducted using mRNA prepared from human B Lymphoma cell line RAMOS as a template with the above pair of primers according to ~~the~~a previous report (J. Biol. Chem., 274:18470-18476, 1999) ~~by~~following the standard method. ~~A~~The nucleotide sequence of the ~~obtained~~resulting PCR product was determined and cDNA encoding a full length human AID protein was obtained (cDNA sequence: SEQ ID NO:7, and amino acid sequence: SEQ ID NO:8).

20 [00332] ~~As a result~~The results indicate that the, human AID protein (SEQ ID NO:8) has extremely high homology in amino acid sequences with ~~a~~the mouse AID protein (SEQ ID NO:2) (FIG. 22). ~~A~~The amino acid sequences in the Cytidine and deoxycytidilate deaminase zinc-binding region which is an active region in the AID protein (both mouse AID and human AID amino acid residues 56 to 94 of SEQ ID NOs: 2 and 8, respectively~~SEQ ID NO:56 to 94~~) were completely consistent (conserved) between mouse and human.

30 [00333] As ~~a~~the partial amino acid sequence (amino acid NO: 116 to 132 in SEQ ID NO:2) of mouse AID protein used for the preparation of anti-AID protein antibody (Example 5) was completely consistent with ~~a~~the corresponding amino acid sequence (amino acid NO:



116 to 132 in SEQ ID NO:8) of human AID protein, the anti-AID protein antibody was expected to ~~comprise cross-reactivity~~ not only with mouse AID protein but also with human AID protein.

[00334] Human AID cDNA obtained in the above was reconstructed according to the standard method in the manner of genetic engineering so that His-AID fusion protein, added with His-tag (a peptide of histidine repeated 10 times) added at the N-terminus of ~~at in the~~ human AID protein, was produced, and an expression vector was prepared by inserting the cDNA into a plasmid pEF-BOS (Unexamined published Japanese patent No. Hei 2-242687). The vector was introduced into a monkey kidney-derived ~~derived~~ cell line, COS7, by lipofection using LIPOFECTAMINE (GIBCO BRL) according to the standard method. The ~~obtained resulting~~ transgenic cells were cultured by ~~the standard methods~~ and His-human AID fusion protein was transiently expressed. His-human AID fusion protein was extracted and purified in the same method as ~~the previously~~ reported, and the production of His-human AID fusion protein was analyzed by Western blotting with the anti-AID antibody prepared in Example 5 and a commercial anti-His tag antibody according to the standard method. As a result, ~~The~~ His-AID protein was detected as a band comprising about 31 kDa molecular weight in all cases using any either antibody.

[00335] Example 17: Determination of exons in the genomic DNA encoding the human AID protein

[00336] Based on the information for the nucleotide sequence of the cDNA encoding the full length human AID protein above, exons in the nucleotide sequences for genomic DNA encoding the human AID protein in the above were determined.

[00337] As a result, ~~it the~~ the genomic DNA was confirmed to consist of 5 exons.

Exon 1 (Nucleotide sequence: SEQ ID NO:11);

Exon 2 (Nucleotide sequence: SEQ ID NO:12);

Exon 3 (Nucleotide sequence: SEQ ID NO:13);

Exon 4 (Nucleotide sequence: SEQ ID NO:14); and

Exon 5 (Nucleotide sequence: SEQ ID NO:15).

[00338] ~~The e~~Exon 1 contains a translation initiation codon ATG which codes for the first methionine (Amino acid No: 1 of SEQ ID NO:8) in the human AID protein; ~~and the~~ initiation codon corresponds to nucleotides NOs: 80 to 82 in SEQ NO: 11.

[00339] Specifically, the genomic DNA including DNA encoding the human AID obtained in the above Examples (SEQ ID NO:9, SEQ ID NO:10 and SEQ ID NO:35) consists of introns and exons as described below and comprises a full length of about 11 kb. FIG 23 schematically shows the structure.

5 <SEQ ID NO:9>

Intron: Nucleotide No $\Theta$ s $\div$  from 1 to 1031

Exon 1: Nucleotide No $\Theta$ s $\div$  from 1032 to 1118

Intron: Nucleotide No $\Theta$ s $\div$  from 1119 to 5514

10 <SEQ ID NO:10>

Intron: Nucleotide No $\Theta$ s $\div$  from 1 to 1064

Exon 2: Nucleotide No $\Theta$ s $\div$  from 1065 to 1212

Intron: Nucleotide No $\Theta$ s $\div$  from 1213 to 2591

Exon 3: Nucleotide No $\Theta$ s $\div$  from 2592 to 2862

15 Intron: Nucleotide No $\Theta$ s $\div$  from 2863 to 3155

Exon 4: Nucleotide No $\Theta$ s $\div$  from 3156 to 3271

Intron: Nucleotide No $\Theta$ s $\div$  from 3272 to 3740

Exon 5: Nucleotide No $\Theta$ s $\div$  from 3741 to 5912

Intron: Nucleotide No $\Theta$ s $\div$  from 5913 to 6564

20 <SEQ ID NO:35>

Intron: Nucleotide No $\Theta$ s $\div$  from 1 to 441

Exon 1: Nucleotide Ne $\Theta$ s $\div$  from 442 to 528

Intron: Nucleotide No $\Theta$ s $\div$  from 529 to 6279

25 Exon 2: Nucleotide No $\Theta$ s $\div$  from 6280 to 6427

Intron: Nucleotide No $\Theta$ s $\div$  from 6428 to 7806

Exon 3: Nucleotide No $\Theta$ s $\div$  from 7807 to 8077

Intron: Nucleotide No $\Theta$ s $\div$  from 8078 to 8370

Exon 4: Nucleotide No $\Theta$ s $\div$  from 8371 to 8486

30 Intron: Nucleotide No $\Theta$ s $\div$  from 8487 to 8955

Exon 5: Nucleotide No $\Theta$ s $\div$  from 8956 to 11067

[00340] Example 18: Amplification of a given partial nucleotide sequence of genomic DNA encoding the human AID protein by PCR, and diagnosis ~~for of~~ the presence or absence  
5 of mutations in the partial nucleotide sequence.

[00341] The AID protein of the present invention may be involved in ~~sideration of the~~  
development of various immunodeficiency and allergic diseases. For example, a given  
immunodeficiency or allergic disease may be caused by mutation or deletion in the  
nucleotide sequence of genomic DNA (especially in an exon) encoding an AID protein.

10 [00342] The presence or absence of such a mutation or deletion in the genomic DNA  
can be analyzed by, for example, the following examples.

(1) A pair of primers comprising a nucleotide sequence complementary to  
a given partial nucleotide sequence of genomic DNA encoding the AID protein in the  
present invention is prepared.

15 (2) Using genomic DNA encoding the AID protein obtained from tissues  
or cells of a patient suffering from immunodeficiency or allergic disease as a  
template, an objective partial nucleotide sequence of the genomic DNA is amplified  
with the pair of primers ~~DNA~~.

(3) By analyzing the presence or absence of a PCR product and a  
20 nucleotide sequence of the PCR product, and comparing the nucleotide sequence with  
a corresponding nucleotide sequence in genomic DNA encoding the AID protein  
derived from a normal person, a mutation or deletion in the genomic DNA is  
identified.

[00343] Specifically, this method enables, for example, not only ~~elucidate~~ elucidation  
25 of the relationship between immunodeficiency or allergic disease and the AID protein, but  
also ~~diagnose the of~~ diseases by the above method in the case that AID protein is a cause of  
~~sideration development~~ of a given type of disease (for example, immunodeficiency or allergic  
disease).

[00344] For the above purpose, the following 15 kinds of primers were designed and  
30 prepared based on a given partial nucleotide sequence in the genomic DNA encoding the  
human AID protein.

Primer: p3 (SEQ ID No. ~~NO~~:20)  
 Primer: p9 (SEQ ID No. ~~NO~~:21)  
 Primer: p10 (SEQ ID No. ~~NO~~:22)  
 Primer: p12 (SEQ ID No. ~~NO~~:23)  
 5 Primer: p14 (SEQ ID No. ~~NO~~:24)  
 Primer: p16 (SEQ ID No. ~~NO~~:25)  
 Primer: p17 (SEQ ID No. ~~NO~~:26)  
 Primer: p19 (SEQ ID No. ~~NO~~:27)  
 Primer: p26 (SEQ ID No. ~~NO~~:28)  
 10 Primer: p29 (SEQ ID No. ~~NO~~:29)  
 Primer: p36 (SEQ ID No. ~~NO~~:30)  
 Primer: p48 (SEQ ID No. ~~NO~~:31)  
 Primer: p59 (SEQ ID No. ~~NO~~:32)  
 Primer: p85 (SEQ ID No. ~~NO~~:33)  
 15 Primer: p86 (SEQ ID No. ~~NO~~:34)

[00345] By PCR using the above primers as ~~in~~ a pairs ~~in~~ of primers by the following combinations, and a genomic DNA isolated from human B lymphoma cell RAMOS as a template, a partial nucleotide sequence coding each target human AID protein was amplified.  
 20 FIG. 21 shows the relative locations of genomic DNA partial nucleotide sequences amplified by each primer pair.

- (1) DNA comprising the nucleotide sequence of SEQ ID NO:31 and DNA comprising the nucleotide sequence of SEQ ID NO:32;
- (2) DNA comprising the nucleotide sequence of SEQ ID NO:20 and DNA  
 25 comprising the nucleotide sequence of SEQ ID NO:22;
- (3) DNA comprising the nucleotide sequence of SEQ ID NO:21 and DNA comprising the nucleotide sequence of SEQ ID NO:30;
- (4) DNA comprising the nucleotide sequence of SEQ ID NO:24 and DNA comprising the nucleotide sequence of SEQ ID NO:25;
- 30 (5) DNA comprising the nucleotide sequence of SEQ ID NO:23 and DNA comprising the nucleotide sequence of SEQ ID NO:27;

(6) DNA comprising the nucleotide sequence of SEQ ID NO:23 and DNA comprising the nucleotide sequence of SEQ ID NO:28;

(7) DNA comprising the nucleotide sequence of SEQ ID NO:23 and DNA comprising the nucleotide sequence of SEQ ID NO:29;

5 (8) DNA comprising the nucleotide sequence of SEQ ID NO:26 and DNA comprising the nucleotide sequence of SEQ ID NO:27;

(9) DNA comprising the nucleotide sequence of SEQ ID NO:26 and DNA comprising the nucleotide sequence of SEQ ID NO:28;

10 (10) DNA comprising the nucleotide sequence of SEQ ID NO:26 and DNA comprising the nucleotide sequence of SEQ ID NO:29;

(11) DNA comprising the nucleotide sequence of SEQ ID NO:34 and DNA comprising the nucleotide sequence of SEQ ID NO:28;

(12) DNA comprising the nucleotide sequence of SEQ ID NO:34 and DNA comprising the nucleotide sequence of SEQ ID NO:29;

15 (13) DNA comprising the nucleotide sequence of SEQ ID NO:33 and DNA comprising the nucleotide sequence of SEQ ID NO:29; or,

(14) DNA comprising the nucleotide sequence of SEQ ID NO:18 and DNA comprising the nucleotide sequence of SEQ ID NO:19;

[00346] The condition for PCR was set by the following manner.

20 [00347] <Reaction solution>

[00348] A total amount of 20.2  $\mu$ l consisting of DDW (8  $\mu$ l), 10 X buffer (2  $\mu$ l), dNTP (2.5 mM each, 2  $\mu$ l), 2 mM primer 1 (2  $\mu$ l), 2  $\mu$ M primer 2 (2  $\mu$ l), genomic DNA isolated from human B Lymphoma cells (185 ng/ $\mu$ l) and Taq polymerase (5 U/ml, 0.2  $\mu$ l), Ex Taq (TAKARA), or Ampli Taq (Perkin Elmer).

25

[00349] <Reaction>

[00350] Reaction was conducted ~~by~~ according to the following conditions (A) or (B).

(A) Conducting 1 cycle of {[reaction at 94°C for 30 sec]} and 40 cycles of {[reaction at 94°C for 10 sec, reaction at 54°C for 30 sec, and reaction at 72°C for 3 min and 30 sec]}, and subsequently stored at 4°C

30

(B) Conducting 1 cycle of {[reaction at 94°C for 30 sec]} and 40 cycles of [{reaction at 94°C for 10 sec, reaction at 55°C for 30 sec, and reaction at 72°C for 2 min and 10 sec},] and subsequently stored at 4°C.

5 [00351] <PCR equipment>

[00352] A commercial PCR device (Perkin Elmer Thermal Cycler 9700 type) was used.

[00353] Example 19: The expression of human AID mRNA in various human organ  
10 tissues

[00354] The expression of human AID mRNA in various human organ tissues was analyzed by RT-PCR according to the standard method (Immunity, 9:1-10, 1998).

[00355] RT-PCR was conducted ~~by using various tissues set in the~~ a human tissue cDNA panel (CLONTECH) as a template according to ~~the~~ standard methods.

15 [00356] AID cDNA was amplified ~~by using~~ primers p17 (SEQ ID NO:26) and p26 (SEQ ID NO:28) prepared ~~in the~~ as above, and Taq polymerase.

[00357] As a control, RT-PCR was conducted in the same manner ~~was conducted~~ using cDNA of G3PDH as a template and GF primer and GR primer (Immunity, 9:1-10, 1998).

20 [00358] FIG. 24 shows the results. As a result, Specific expression of mRNA was confirmed in lymph node and tonsil. This result was consistent with the experimental result in which the expression of mRNA for mouse AID was observed in the various lymphatic tissues (Examples 8 and 9).

[00359] On the other hand, when RT-PCR was conducted with ~~the~~ a saturated cycle  
25 number in the same manner as the above in the above RT-PCR, the expression of AID mRNA was observed in almost all analyzed organs.

[00360] Example 20: Localization of human AID gene on human chromosomes

[00361] Localization of the human AID gene on human chromosomes was analyzed  
30 by ~~Fluorescence~~ fluorescence in situ hybridization (FISH) method (Experimental Medicine, Suppl. "Genetic Engineering Hand Book" published by Yodosha, 1992, p. 271-277).

[00362] Genomic DNA including human AID gene (exon 1 to exon 5), isolated in the above, which was labeled with biotin-11-dUTP (Sigma) by the nick translation method, was used as a probe for hybridization.

[00363] The probe was hybridized with chromosomes in metaphase human cells.

5 Hybridization signals were detected using fluorescein isothiocyanate-avidin (DCS) (Vector Laboratories).

[00364] FIG. 25 shows the results. ~~As a result,~~The human AID gene was found to be localized on chromosome 12p13. ~~Moreover, this location was revealed to be~~is near 12p13.1 which is the location for ~~the above-APOBEC-1 comprising which has~~ a relatively high amino acid sequence homology with the AID protein, and has the same cytidine deaminase activity as ~~that of~~ the AID protein.

[00365] It has been reported that some abnormality on human chromosome locus 12p13.3-12p11.2, 12p13.2-12p24.1 and 12p13 may be involved in diseases such as acrocallosal syndrome, inflammatory bowel syndrome, familial periodic fever, respectively, however, the causative gene thereof has not been traced yet. ~~It has been suggested that~~ The human AID gene of the present invention may be involved in ~~sideration~~ the development of such diseases.

[00366] All references and patents cited herein are incorporated by reference in their entirety.

20

## INDUSTRIAL APPLICABILITY

[00367] The AID protein of the present invention can be considered to have a function of regulating various biological mechanisms required for generation of antigen-specific immunoglobulins (specific antibodies), which eliminate non-self antigen (foreign antigen, self-reacting cells, etc.) that triggers various diseases. More specifically, the AID protein of the present invention can be considered to be one of the enzymes that plays an important role in ~~the~~ genetic editing such as RNA editing, and so on occurring in germinal center B cells, such as the activation of B cells, class switch recombination of immunoglobulin genes, somatic hypermutation, and affinity maturation, which are the mechanisms for generation of immunoglobulins having high specificity ~~to~~ for antigens.

30

[00368] The dysfunction of the AID protein of the present invention can be the cause for the of humoral immunodeficiency since it induces failure of germinal center B cell function, such as antigen-specific B cell activation, class switch recombination, and somatic mutation. ~~Reversely~~Conversely, the breakdown of the regulation of AID protein may induce allergy allergic diseases or autoimmune diseases since it can cause inappropriate B cell activation and needless class switch recombination and somatic mutation.

[00369] Therefore, regulation of the function of AID protein and the gene encoding it enables preventiong and treatmenting of various immunodeficiencies, autoimmune diseases, and allergies, which result from, for example, B cell dysfunctions (e.g. IgA deficiency, IgA nephropathy,  $\gamma$  globulinemia, hyper IgM syndrome, etc.) or class switch deficiency of immunoglobulin. Thus, the AID protein and the gene encoding the AID protein can be targets for the development of drugs for therapy of the diseases mentioned above.

[00370] Examples of diseases whose onset prevention, symptom remission, therapy and/or symptomatic treatment effect is expected by regulating the function of the AID protein of the present invention or the gene encoding it include, for example, primary immunodeficiency syndrome with congenital disorder of immune system, mainly immunodeficiencyies considered to develop by B cell deficiency, decrease, or dysfunction (e.g. sex-linked agammaglobulinemia, sex-linked agammaglobulinemia with growth hormone deficiency, immunoglobulin deficiency with high IgM levels, selective IgM deficiency, selective IgE deficiency, immunoglobulin heavy chain gene deletion,  $\kappa$  chain deficiency, IgA deficiency, IgG subclass selective deficiency, CVID (common variable immunodeficiency), infantile transient dysgammaglobulinemia, Rosen syndrome, severe combined immunodeficiency (sex-linked, autosomal recessive), ADA (adenosine deaminase) deficiency, PNP (purine nucleoside phosphorylase) deficiency, MHC class II deficiency, reticular dysplasia, Wiskott-Aldrich syndrome, ataxia telangiectasia, DiGeorge syndrome, chromosomal aberration, familial Ig hypermetabolism, hyper IgE syndrome, Gitlin syndrome, Nezelof syndrome, Good syndrome, osteodystrophy, transcobalamin syndrome, secretary bead syndrome, etc.), various diseases with antibody production deficiency that are secondary immunodeficiency syndromes with a disorder of the immune system caused by an acquired etiology (for example, AIDS, etc.), and/or various allergic diseases (e.g., bronchial asthma, atopic dermatitis, conjunctivitis, allergic rhinitis, allergic enteritis, drug-induced



allergy, food allergy, allergic urticaria, glomerulonephritis, etc.). These could be targets for drug development.

[00371] Namely, the AID protein of the present invention, a fragment thereof, a DNA encoding the AID protein, a fragment thereof, and an antibody against the AID protein are  
5 useful as reagents for developing drugs for prevention and therapy of such diseases.

[00372] Also, the DNA itself is useful as an antisense drug regulating the function of the AID gene at a~~the~~ gene level, and in a~~use in~~ gene therapy. The protein, or the fragments thereof (e.g. enzyme active site), itself is useful as a drug.

[00373] Furthermore, an antibody reactive to the AID protein of the present invention  
10 or a fragment thereof is extremely useful as an antibody drug ~~by~~to regulating~~regulate~~ functions of the AID protein.

[00374] Furthermore, the gene (DNA), protein, and antibody of the present invention are useful as reagents for searching for substrates (e.g., RNA, etc.) that ~~interacting~~ (binding) with the protein (enzyme) of the present invention, or other auxiliary proteins associated with  
15 the protein of the present invention, and for developing drugs targeting the substrates and auxiliary proteins.

[00375] Furthermore, a method for identifying a substance that regulates production of the AID protein of the present invention, or transcription of a gene encoding the AID protein into mRNA ~~are~~is extremely useful as a means to develop drugs for therapy and prevention of  
20 various diseases (especially, immunodeficiency and allergic disease), in which the above-mentioned AID protein or AID gene is considered to be involved.

## ABSTRACT

Genes encoding novel proteins named AID (Activation-Induced cytidine Deaminase), ~~having that are~~ structurally related ~~relationship~~ to APOBEC-1, ~~one of an~~ RNA editing enzymes, and ~~having~~ a cytidine deaminase activity similar to APOBEC-1, ~~have~~ been found by  
5 preparing cDNA libraries ~~for~~ from mouse B cell clone CH12F3-2; (which undergoes class switch recombination from IgM to IgA at an extremely high rate ~~together with~~ after activation of the cells by stimulation with cytokines), with and without stimulation ~~ng~~ with cytokines, and performing subtraction cloning using the libraries.